

A global climatology of sting-jet extratropical cyclones

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Gray, S. L. ORCID: <https://orcid.org/0000-0001-8658-362X>,
Volonté, A. ORCID: <https://orcid.org/0000-0003-0278-952X>,
Martinez-Alvarado, O. ORCID: <https://orcid.org/0000-0002-5285-0379> and Harvey, B. J. ORCID: <https://orcid.org/0000-0002-6510-8181> (2024) A global climatology of sting-jet extratropical cyclones. *Weather and Climate Dynamics*, 5 (4). pp. 1523-1544. ISSN 2698-4024 doi: <https://doi.org/10.5194/wcd-5-1523-2024> Available at <https://centaur.reading.ac.uk/119944/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.5194/wcd-5-1523-2024>

Publisher: EGU

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



A global climatology of sting-jet extratropical cyclones

Suzanne L. Gray¹, Ambrogio Volonté^{1,2}, Oscar Martínez-Alvarado^{1,2}, and Ben J. Harvey^{1,2}

¹Department of Meteorology, University of Reading, Reading, UK

²National Centre for Atmospheric Science, University of Reading, Reading, UK

Correspondence: Suzanne L. Gray (s.l.gray@reading.ac.uk)

Received: 15 May 2024 – Discussion started: 28 May 2024

Revised: 18 October 2024 – Accepted: 24 October 2024 – Published: 17 December 2024

Abstract. Sting jets have been identified in the most damaging extratropical cyclones impacting northwest Europe. Unlike the cold conveyor belt and other long-lived cyclone wind jets, sting jets can lead to regions of exceptionally strong near-surface winds and damaging gusts over just a few hours and with much smaller wind “footprints”. They descend into the frontal-fracture region found in warm-seclusion cyclones. Previous research has focused almost exclusively on North Atlantic–European cyclones, but there are no known physical reasons why sting jets should not develop elsewhere, and recognition of their existence can inform weather nowcasting and wind warnings. We have produced the first climatology of sting-jet cyclones over the major ocean basins. A sting-jet precursor diagnostic has been applied to more than 10 000 warm-seclusion cyclones in the top intensity decile, tracked using 43 extended winters of ERA5 reanalysis data. Cyclones with sting-jet precursors are found to occur over the North Pacific and Southern oceans for the first time, and they are more prevalent in the Northern Hemisphere (27 % of all top decile cyclones) compared to the Southern Hemisphere (15 %). These cyclones have distinct characteristics to those without the precursor, including initiating closer to the Equator, deepening faster in mean-sea-level pressure (MSLP), and having stronger near-surface winds, even in the reanalysis data that are too coarse to fully resolve sting jets. Composite analysis reveals systematic differences in structural evolution, including in potential vorticity (PV) and jet crossing. These differences evidence the climatological consequences of strong diabatic cloud processes on cyclone characteristics, implying that sting jets are likely to be enhanced by climate change.

1 Introduction

Sting jets have been identified in several of the most damaging extratropical cyclones impacting northwest Europe (see the review paper by Clark and Gray, 2018) since they were first identified in 2004 from a re-examination of observations of the 1987 Great October Storm that devastated southern England (Browning, 2004; Clark et al., 2005). As a recent example, Volonté et al. (2023a, b) present evidence of the sting jets in storm Eunice, which caused extensive wind damage in southern England in February 2022. The possibility of sting jets enhancing damaging winds and gusts is now routinely considered by the (UK) Met Office in forecasting and generating weather warnings, and the term sting jet is used in their communications and by the British media. The aim of this study is to identify and characterise sting-jet cyclones and their distinctiveness from non-sting-jet cyclones in the three major ocean basins. Recognition of their potential existence is a first step towards improved nowcasting and warning of associated wind risk. Sting jets are also a consideration for the currently uncertain projections of climate change effects on cyclone wind intensity (Catto et al., 2019).

Strong near-surface winds in extratropical cyclones are mainly associated with one of two major airflows termed conveyor belts: the warm conveyor belt ahead of the cold front and the cold conveyor belt poleward of the cyclone centre that travels rearwards relative to the cyclone motion. Their definitions, and those of other major cyclone features, date back to at least the 1980s (see Browning and Roberts, 1994), and the associated surface winds typically last many hours, yielding wind speed footprints up to hundreds of kilometres wide by thousands of kilometres long. In contrast, sting jets are, as defined by Clark and Gray (2018), coherent airflows that descend over just a few hours from within the cloud

heads visible in intense cyclones. They are distinct from the conveyor belt jets, although they can interact with the cold conveyor belt jet, and can lead to regions of exceptionally strong near-surface winds and particularly damaging gusts, with much smaller footprints (50–100 km across; Hewson and Neu, 2015). Their small-scale, transient nature is one reason for their much more recent identification: they can pass between operational surface weather stations, although they have occasionally been captured by wind profilers (Volonté et al., 2018; Parton et al., 2009) and Doppler radar (Browning et al., 2015), and their full representation by numerical models requires the latest high-resolution weather forecast models. Sting jets descend into the frontal-fracture region of cyclones, developing according to the Shapiro–Keyser conceptual model (Shapiro and Keyser, 1990). One distinctive aspect of Shapiro–Keyser cyclones is that a seclusion of warm air forms in their centre as they approach maturity, which is delimited by a bent-back warm front wrapping around the poleward side of the low-pressure centre; this structure contrasts with the occluded front characteristic of cyclones that develop according to the other major conceptual model, the Norwegian model (Bjerknes, 1919).

Since their identification, knowledge of the characteristics and physical mechanisms causing sting jets has advanced considerably. Analysis of cyclone case studies and idealised model simulations led to the development of a conceptual model of sting-jet evolution, revealing the role of mesoscale instabilities, including conditional symmetric instability (CSI), in their generation, descent, and acceleration (Gray et al., 2011; Volonté et al., 2020, 2018). A climatological analysis, using a precursor diagnostic based on the potential for CSI release by descent in the cloud head, showed that sting jets exist in perhaps one in three North Atlantic cyclones, particularly explosive ones (Hart et al., 2017). Sting-jet frequency is projected to increase in the future (Martínez-Alvarado et al., 2018; Manning et al., 2021, 2023). However, all this research has focused exclusively on cyclones crossing the North Atlantic and affecting northwest Europe. A single case of sting-jet activity in a cyclone originating in the Mediterranean has recently been published (Brâncuş et al., 2019), but no published cases conclusively diagnosing sting-jet existence exist for other extratropical regions, and there is no published climatology beyond the North Atlantic. Despite this, there are no known physical reasons why sting jets should not develop over other extratropical oceanic basins.

In developing this climatology, we address two sets of questions related to their locations and structures.

1. Is the sting jet a global extratropical phenomenon or is it limited to North Atlantic cyclones? What are the tracks and prevalence of these cyclones over the major ocean basins and how do they differ from those of non-sting-jet cyclones?
2. What are the structural differences between sting-jet and non-sting-jet cyclones globally and in regional sub-

sets? For example, are sting-jet cyclones more intense, as is found for cyclones crossing the North Atlantic (Hart et al., 2017)?

Unambiguous identification of sting jets in cyclone case studies requires detailed analysis of numerical weather prediction model output from a model with high enough resolution to resolve the sting jet: Clark and Gray (2018) conclude that a minimum horizontal grid spacing of 10–15 km is required, with correspondingly high enough vertical grid spacing to resolve the slantwise motions. The analysis ideally includes analysis of air parcel trajectories to disentangle the low-level wind jets and observational analysis to support the model prediction. Thus, generation of a climatology requires an alternative approach, both due to the challenge of identifying sting jets and because such analysis typically makes use of long-term reanalysis products or climate model outputs that are produced using models that have insufficient resolution to resolve sting jets. Rather than unambiguously identifying sting jets in cyclones, instead cyclones that are likely to contain sting jets can be identified.

Here a sting-jet precursor (SJP) diagnostic based on the presence of CSI within the cloud head, developed by Martínez-Alvarado et al. (2013) and previously used to produce the first North Atlantic climatologies of sting-jet cyclones (Martínez-Alvarado et al., 2012; Hart et al., 2017), has been adapted and applied to tracked cyclones with warm seclusions in 43 years of extended-winter ERA5 reanalysis data across the entirety of the extratropics in both hemispheres. In addition to production of the first global climatology of extratropical sting-jet cyclones, novel aspects of this study include the development of a new approach to identify cyclones containing warm seclusions (Shapiro–Keyser cyclones) using an image segmentation approach; the development of a set of 33 notable storms, used to refine the SJP diagnostic, holistically assessed for likely sting-jet presence (using observations, the previous literature, and detailed analysis of the location and timing of the regions with CSI); and comprehensive comparison of the characteristics and composite structure of cyclones with and without the SJP, mainly through the use of ERA5 data but also using an observational product: a blend of scatterometer observations with bias-corrected ERA5 data.

The remainder of this paper is structured as follows. The reanalysis data and cyclone-tracking algorithm, observational data, and compositing method used are described in Sect. 2, followed by the warm-seclusion algorithm and SJP diagnostic in Sect. 3. The results section (Sect. 4) begins with a short analysis of how the proportion of warm-seclusion cyclones varies with cyclone intensity before a detailed analysis of how the frequency, spatial and temporal variability, intensity metrics (including from the blended scatterometer data), and composite structure of cyclones with and without the SJP compare. The conclusions section (Sect. 5) finishes the paper.

2 Methods I: datasets, cyclone tracking, and compositing

2.1 Reanalysis data and cyclone tracking

Cyclones were identified using the ERA5 dataset, which is the latest comprehensive global reanalysis dataset produced by the European Centre for Medium-Range Weather Forecasts (ECMWF). ERA5 was produced using cycle Cy41r2 of ECMWF's integrated forecast system (IFS), which was operational from 8 March to 21 November 2016. In this cycle the IFS model had a horizontal spectral truncation of TL639 (equivalent to 31 km grid spacing) with 137 terrain-following hybrid-pressure levels up to a height of 80 km. Atmospheric data are available on a regular latitude–longitude grid with 0.25° (~ 30 km) spacing. For this study, cyclones were identified and tracked using hourly ERA5 data for 43 years of extratropical (defined as 40° – 80°) extended winters in the Northern Hemisphere (October–March) from 1979–1980 to 2021–2022 and in the Southern Hemisphere (April–September) from 1980 to 2022.

The TRACK algorithm (version 1.5.2) was used to track the cyclones (Hodges, 1995, 1999, 2023). The cyclones were identified as maxima of spatially smoothed 850 hPa relative vorticity (ξ_{850}) fields, with the smoothing performed using spectral filtering to retain features with total wave numbers between T5 and T63 (where T refers to the triangular truncation of the spherical harmonics), which equates to an approximately smallest-resolved scale of 320 km for circular features. Cyclone features with maximum smoothed ξ_{850} values exceeding 10^{-5} s^{-1} were retained and tracked with the tracks retained if they lasted for more than 1 d with a total track distance exceeding 500 km; this duration was chosen to retain all of the cyclones in the notable storms list used for verification. Spatial statistics such as track and genesis density were computed from the tracks using spherical kernels (Hodges, 1996).

After calculating the tracks using the hourly ERA5 data (as higher-temporal-resolution data improves the track accuracy), the temporal resolution of the track files was then degraded to the 6-hourly times (00:00, 06:00, 12:00, and 18:00 UTC only) at which the SJP was evaluated. To keep the number of tracks manageable for further processing, we choose to focus on the top decile of tracks in terms of the maximum intensity (defined as the maximum value of smoothed ξ_{850}) along the extratropical portion of the track and to consider only cyclones in which a warm seclusion is present at the time of maximum intensity (as sting jets have only been identified as descending towards the surface in such Shapiro–Keyser cyclones). Percentiles were computed separately for each hemisphere, and the 90th percentile values for spatially smoothed ξ_{850} are 12.72×10^{-5} and $13.95 \times 10^{-5} \text{ s}^{-1}$ in the Northern Hemisphere and Southern Hemisphere, respectively. Note that the Northern Hemisphere absolute threshold values were also used for the North

Atlantic and North Pacific regions. These thresholds equate to local percentiles of 86 % for both regions, implying that cyclones in these basins are typically more intense than in the Northern Hemisphere overall. After sub-selecting cyclones in which a warm seclusion was diagnosed (as described in Sect. 3.1), the SJP diagnostic (Sect. 3.2) was applied to sets of 5550 Northern Hemisphere and 5207 Southern Hemisphere cyclones. For each cyclone, the fields required for the SJP diagnostic were extracted from the ERA5 dataset in a square domain of $\pm 15^\circ$ in both latitude and longitude (with a spherical cap projection) from each track point from 42 h before to 42 h after the time of maximum intensity. To enable comparison with previous studies that considered all cyclones in the North Atlantic, a randomly selected year of data was also processed for all cyclones in that year.

2.2 Cyclone composites

The structures of the cyclones with and without the SJP (excluding marginal cases) have been compared through compositing cyclone fields according to the time relative to the time of maximum cyclone intensity and the ocean basin (Northern or Southern Hemisphere, North Atlantic, or North Pacific). The North Atlantic and North Pacific basins are defined throughout this study as being between the longitudes of 80° W – 30° E and 120° E – 120° W , respectively¹. The cyclones are rotated prior to compositing such that their travel direction points eastward, with the aim of better aligning the major cyclone features, such as fronts, between the cyclones. Cyclones with atypical travel directions, defined as in Sect. 3.2, are excluded from the composites (at the time that the direction is anomalous only), as after rotation the orientation of their major features tends to be anomalous.

Cyclone relative fields are extracted for the SJP calculation; gridded using a rotated-pole coordinate system centred on the MSLP centre of each cyclone; and an additional field, potential vorticity (PV), was extracted in the same way for compositing. As described in Sect. 2.1, the extracted domain is a square that extends $\pm 15^\circ$ from the storm centre. After compositing, the domain was reduced to $\pm 12^\circ$ from the storm centre to exclude the outer portions of the domain where the fields were more noticeably noisy due to missing data for some of the cyclones after rotation; however note that there will still be slightly fewer data points in the composites near the edges of the domain shown, especially in the corners.

2.3 Observational data

We utilise several independent sources of satellite data in this study, as input into the expert judgement evaluation process

¹For compositing, the location of the tracked vorticity centre was considered for convenience rather than its associated mean-sea-level pressure (MSLP) centre (as used in the rest of the analysis)

for the set of notable storms outlined in Sect. 3 and to compare the sets of cyclones with and without a diagnosed SJP.

Passive radiometer images were obtained from three sources: the High Rate Spinning Enhanced Visible Infra-Red Imager (SEVIRI) IR10.8 band from the Meteosat Second Generation (MSG) 0° satellite, which covers the Atlantic sector and is available at 15 min frequency from 2004–present; the Advanced Very High Resolution Radiometer (AVHRR) IR10.8 band from the METOP (Meteorological Operational) satellites, which are polar orbiters and therefore available on daily swaths covering limited areas; and the Geostationary Ring IR10.8 Multimission Image produced by EUMETSAT, which merges 3-hourly images from several geostationary satellites to provide global coverage. In each case, we use the satellite image closest in time to that of the maximum intensity of the cyclone, favouring the higher-resolution AVHRR images over the SEVIRI/Multimission images when they are available.

Ocean surface wind data are obtained from the Cross-Calibrated Multiplatform Version 3.0 dataset (CCMP3.0; Mears et al., 2022a), which provides gridded vector 10 m wind data over ocean points on a $0.25^\circ \times 0.25^\circ$ grid from 1993 to 2019. CCMP3.0 uses a variational analysis method to blend ocean surface wind retrievals from a wide range of satellite microwave sensors with a background field of adjusted 10 m neutral winds from ERA5 to produce a regular 6-hourly dataset (Atlas et al., 2011). The ERA5 adjustment involves removing ocean currents vectors to obtain wind speeds relative to the ocean surface and applying a quantile-mapping technique to bias correct strong wind values, which are typically too weak in ERA5.

3 Methods II: identification of cyclones with sting-jet precursors

A set of 33 notable storms was used to test and refine the warm-seclusion and SJP algorithms described in the following two subsections before applying them to the complete cyclone datasets. These notable storms were intense cyclones generally associated with severe impacts, which in some cases were the subject of published case studies. They were all identified as $>$ the 90th percentile in intensity warm-seclusion cyclones (see Sect. 3.1) and included cases from the North Atlantic (both near the North American east coast and further east all the way to western Europe), North Pacific (including the North American west coast), and Southern Ocean basins. The method of assessing the notable storms is described in the Supplement (Sect. S2) together with lists of the storms in Tables S1 and S2.

3.1 Identification of warm-seclusion cyclones

The maps in Fig. 1 illustrate the algorithm used for warm-seclusion cyclone identification. Three examples are shown:

two cyclones identified as having a warm seclusion (one from each hemisphere) and one that failed the identification test. (Note that the two warm-seclusion cyclones are in the notable storms set.) The two panels shown for each cyclone show the low-level cyclone structure (the MSLP field with the identified cyclone core together with the 850 hPa wet-bulb potential temperature (θ_w) field) and warm-seclusion diagnostic process, respectively. A short description of the algorithm follows.

Warm-seclusion identification is only applied to a track if the associated cyclone centre enters the midlatitudes (i.e. the MSLP cyclone centre associated with the tracked vorticity feature is, for at least one 6-hourly time step, within 40° and 80° of latitude in either hemisphere). The time chosen for performing the identification is that of cyclone maximum intensity (when ξ_{850} is at a maximum along the track) while the cyclone centre is in the midlatitudes. The minimum value of MSLP at the time of maximum intensity in a 5° radius (using a geodesic great circle radius, B-spline interpolation, and steepest-descent minimisation) centred on the track point is identified as the cyclone centre. Then, the pressure core is defined as the region where MSLP is less than 4 hPa above its cyclone-centre minimum and less than 300 km away from it.

A type of image segmentation algorithm called a watershed algorithm is then run on the 850 hPa θ_w to identify distinct watersheds within the warm sector (using a footprint size of 8×8 grid points). The warm sector is defined as the region where the 850 hPa θ_w is warmer than its maximum value in the cyclone core minus 2 K. The watershed algorithm works by considering the θ_w values within this warm sector as an inverted local topography (i.e. θ_w maxima are valleys). The algorithm then “floods” basins from markers defined as local maxima in θ_w (lowest points of valleys) until the basins attributed to different markers meet on watershed lines, thus defining the watersheds. If the watershed that overlaps the cyclone pressure core with the most points is all contained within a circle with a radius of 600 km centred at the cyclone centre, the cyclone is identified as having a warm seclusion (provided this watershed exceeds a threshold size of 25 grid points). In the right-hand-side maps in Fig. 1, the watershed boundary is highlighted with a thick green contour if a warm seclusion has been identified. Otherwise, the cyclone is identified as a non-warm-seclusion cyclone, and the watershed boundary is highlighted with a thick red contour. Our warm-seclusion identification method includes several user-defined thresholds. The values for these thresholds were optimised by analysing the behaviour of the algorithm for the set of notable storms as well as other cyclones to provide the best agreement between subjective and algorithm-based warm-seclusion identification. The starting point for our algorithm development was the algorithm used to identify cyclones with warm seclusions by Manning et al. (2021). Their algorithm identifies warm seclusions as localised warm regions, where 850 hPa θ_w is at least 2 K warmer than the sur-




Figure 1. Examples of maps used for evaluation of the warm-seclusion identification algorithm. Left-hand-side maps include MSLP (grey contours, hPa) and 850 hPa θ_w (shading, every 2 K), with warm-sector areas (see definition in the text) highlighted by thick magenta contours. The blue circle has a radius of 300 km and is centred over the cyclone centre. The blue-shaded region covers the cyclone pressure core. Right-hand-side maps include MSLP (grey contours, hPa) and all warm-sector watersheds identified by the algorithm (discrete colours). The thick green or red contour highlights the watershed that overlaps the pressure core with the most points, and its colour depends on whether the cyclone is identified as having a warm seclusion (green) or not (red). The red circle for warm-seclusion identification has a radius of 600 km and is centred over cyclone centre.

roundings, that overlap with an MSLP core defined as points where the MSLP is within 6 hPa of the minimum values. The most important difference to the Manning et al. (2021) approach is that, partly to increase efficiency, we have used a watershed algorithm to identify the warm core rather than identifying a warm peak by searching in turn along each latitudinal row and longitudinal column of the model data. We also used image processing to identify the cyclone MSLP

core, whereas Manning et al. (2021) used the same line-by-line search to identify the cyclone core.

3.2 Sting-jet precursor diagnosis, including calibration and verification across ocean basins

The SJP diagnostic, based on the methodology outlined in Martínez-Alvarado et al. (2013), is applied to the cyclone

data from 12 h before to 6 h after the time of cyclone maximum intensity to focus on the time period when sting jets have been found most likely to occur in case studies. The SJP diagnostic uses a metric for CSI, specifically the energy available to downdraught air through the release of CSI termed the downdraught slantwise convective available potential energy (DSCAPE) (Gray et al., 2011). The CSI assessment is then complemented by a set of additional criteria to ascertain whether the CSI is within an environment conducive to the formation of sting jets. These criteria consist of an evaluation of the availability of moisture for CSI release and the location of CSI relative to the cyclone centre, the bent-back front, and the cloud head. Model grid points meeting the threshold DSCAPE value and environmental criteria are hereafter referred to as CSI points. This methodology has been used, with variations, in climatological assessments (Martínez-Alvarado et al., 2012; Hart et al., 2017), climate change studies (Martínez-Alvarado et al., 2018), and in near-real-time forecasting operations (Gray et al., 2021).

The precursor diagnostic settings used are the same as those in Gray et al. (2021) with the following modifications to the location of CSI. For CSI release to be considered to have the potential to generate sting jets, CSI must be located in the cyclone's cloud head. Therefore, any CSI diagnosed in a sector (of a circle centred on the cyclone) associated with the cyclone's warm conveyor belt must be disregarded. In Gray et al. (2021), the warm conveyor belt sector was defined as an asymmetric sector between 60 and -100° , with these angles measured anticlockwise with respect to the cyclone's direction of travel. Here this sector is symmetric with respect to this direction and is defined by the angle 2α , which was set to 240° , i.e. the sector spans from 120 to -120° .

The cyclone's travel direction is computed at each track point based on the location of the cyclone's centre, using a forward-difference approach for the first time step, a backward-difference approach for the last time step, and a centred-difference approach for any other time step in the cyclone's track. For the forward and backward differencing, the time interval between the chosen track points was 6 h. The centred-difference method was implemented using track points from 12 h before to 12 h after the time of the track point of interest, except for the second and second-to-last time steps when 6 h was used. The longer time interval was preferred when possible because this improved the smoothness of the travel direction's evolution. This implementation differs from Gray et al. (2021), in which centred differencing was used for all time steps, but the first and last time steps were discarded.

The cloud head sector assessment assumes a typical eastward and poleward cyclone travel direction. Therefore, an additional requirement was introduced in this work to flag cyclones with atypical tracks. On inspection, these cyclones were sometimes those close to maturity, tracking slowly with a strong poleward component to their track, and occasionally those earlier in their life cycle performing small track loops,

meaning that for a short period the travel direction had a strong westwards or equatorward component. Measuring the angles with respect to north (such that east is 90°), a cyclone track was considered atypical if the cyclone's travel direction was between 180 and 315° in the Northern Hemisphere and between 225 and 360° in the Southern Hemisphere at any time when the SJP diagnostic was calculated.

To classify the cyclones according to their likelihood of containing a sting jet (in reality), the total number of CSI points diagnosed for each cyclone is considered. The total (rather than, e.g. an instantaneous maximum) is used because there is uncertainty about when CSI appears in a low-resolution model and so the exact timing is not relevant as long as it is around the time of maximum intensity. The cyclones have been classified into three categories according to the output of the SJP diagnostic: sufficient CSI, so likely to produce sting jets (TRUE, > 16 CSI points); insufficient CSI, so unlikely to produce sting jets (FALSE, < 9 CSI points); and intermediate CSI, so neither likely nor unlikely to produce sting jets (MARGINAL, $9 \leq \text{CSI points} \leq 16$). We assessed and refined the methodology using the set of notable storms. Example AVHRR and CCMP images are shown in Fig. 2 for two of the notable storms, illustrating strong 10 m winds ahead (in a cyclonic sense) of the banded cloud head tips. In particular, the results were found to be largely insensitive to the precise values used for α and the time interval used to calculate the cyclone travel direction, and, while varying the threshold numbers of CSI points led to changes in the classification of some cyclones, these changes were found to be relatively small. The output of the sting-jet precursor algorithm is illustrated in Fig. 3, corresponding to the storms in Fig. 2, showing that it is correctly identifying instability in the cloud head tip.

4 Results

4.1 Characteristics of warm-seclusion cyclones

As the SJP diagnostic was only applied to those cyclones exceeding the 90th percentile in maximum intensity and having a diagnosed warm seclusion, the proportion of cyclones with a warm seclusion as a function of the intensity decile is presented in Fig. 4 to show the effect of the warm-seclusion constraint. There is a clear more-than-linear increase in the fraction of cyclones that are identified as having warm seclusions as their maximum intensity increases, with the fraction of warm-seclusion cyclones being similar for both hemispheres and the three major oceanic basins. The fraction of warm-seclusion cyclones is 0.42–0.45 for cyclones in the lowest-intensity decile, increasing to 0.73–0.77 for cyclones in the highest-intensity decile such that the vast majority of cyclones in the top decile are diagnosed as having a warm seclusion. Consequently, the warm-seclusion constraint reduced the number of top-decile cyclones evaluated using the




Figure 2. Satellite observations of the example storms from Fig. 1a and b. Left-hand panels show 10 m wind speed values (units in m s^{-1}) from the CCMP3.0 blended wind product at the times indicated together with contours of MSLP from ERA5 (with a 4 hPa interval), and right-hand panels show IR10.8 band images from the AVHRR instrument for the image that is closest in time to the time shown. In both cases, the data are re-projected onto the same projection used in Fig. 1, but note that the area shown is more zoomed in than in Fig. 1. For the full details of the dataset used, see Sect. 2.3.

SJP diagnostic by less than 30 %, although, due to the computational expense of applying the algorithm, this reduction was still considered worthwhile (and greater benefit would be obtained applying the SJP diagnostic to a broader cyclone intensity range).

4.2 Frequency of cyclone types

The absolute and relative frequencies of cyclones with and without the SJP (termed SJP and nSJP cyclones, respectively) are shown in Table 1. Of the more than 10 000 warm-seclusion cyclones evaluated, 29 % had an SJP and 55 % did not have an SJP (with the remainder categorised as marginal). Assuming that a sting jet would not have been able to descend towards the surface in all the cyclones without a warm seclusion (irrespective of the presence of a sting-jet precursor), 21 % of all top intensity decile cyclones may have had a sting jet. Cyclones in the Southern Hemisphere are only just over half as likely to have the SJP as those in the Northern Hemisphere (20 % of evaluated cyclones compared to 37 %). The percentage of cyclones in each precursor category was similar for North Atlantic and North Pacific basins and when considering the whole Northern Hemisphere.

The percentage of SJP cyclones found for the North Atlantic in this 43-year climatology based on ERA5 data (38 % of evaluated cyclones) is similar to that found by Hart et al. (2017) for this region using 32 years of data from the preceding ERA-Interim reanalysis for the same extended-winter season and using a very similar SJP diagnostic. They found that 32 % of the 5447 cyclones tracked had the precursor; an almost identical percentage (33 %) was also obtained from analysis of a climate model simulation of the present-day climate by Martinez-Alvarado et al. (2018). However, our results are not directly comparable due to differences in the selection of cyclones to evaluate and in the methodology. In particular, Hart et al. (2017) evaluated all cyclones lasting at least 2 d with a total track distance exceeding 1000 km, whereas here only warm-seclusion cyclones in the top intensity decile lasting at least 1 d and with a track distance exceeding 500 km were evaluated.

To assess the consistency between the current and previous results that apply the SJP diagnostic to North Atlantic cyclones and provide indicative results for how the percentage of SJP cyclones varies with the cyclone intensity decile, all tracked cyclones (all intensities and both with and without a warm seclusion) with minimum track durations of both 1




Figure 3. Illustration of the implementation of the SJP diagnostic, showing MSLP (black contours, hPa), 850 hPa θ_w (blue contours, K), and 650 hPa relative humidity with respect to ice above 80 % (grey contour and stippling) at (a) 06:00 UTC on 15 December 2011 and (b) 18:00 UTC on 20 August 2021. The red lines represent cyclone tracks, with the red circles marking the cyclone centres at the validation times. Purple and pink dots represent CSI points satisfying and not satisfying the additional location criteria, respectively.

and 2 d (and associated minimum track distances of 500 and 1000 km, respectively) were evaluated for this ocean basin for a single winter season. The 2011–2012 season chosen is representative of the 43-year winter climatology in terms of the percentages of top intensity decile North Atlantic warm-seclusion cyclones found in the SJP, nSJP, and marginal categories for the same 1 d minimum track duration (Table 1). This season was also included in the period evaluated by Hart

et al. (2017) and contains Cyclone Friedhelm from the set of notable storms. Comparison of the top decile 2011–2012 winter North Atlantic cyclones with all North Atlantic cyclones from that winter (with the same track constraints) reveals that SJP cyclones occur about twice as frequently in the former, providing strong evidence that SJPs are preferentially found in more intense cyclones, consistent with Hart et al. (2017) when comparing all cyclones to explosively develop-

Table 1. Absolute and relative (%) frequency of SJP, nSJP, and marginal SJP warm-seclusion cyclones and non-warm-seclusion cyclones (with minimum 1 d track duration) by hemisphere, major oceanic basin, and overall for the 43-year winter climatology and the 2011–2012 frequencies for the North Atlantic for two choices of minimum track duration (in all cases, the 1 and 2 d minimum track durations are associated with minimum total track distances of 500 and 1000 km, respectively). The top decile intensity thresholds are calculated separately for each hemisphere (but not for each oceanic basin), and consequently, a larger proportion of cyclones are considered in the North Atlantic and North Pacific subregions than in the two hemispheres. The values in brackets are the percentages of all cyclones and of warm-seclusion cyclones only, respectively. Note that the percentages of marginal and non-warm-seclusion cyclones (percentages of the total cyclones) are fairly consistent across regions (11 %–14 % and 23 %–27 %, respectively).

Region	Constraints		Winters	Warm-seclusion cyclones			Non-warm-secl.	Total
	Max ξ_{850}	Length (d)		SJP	nSJP	Marginal		
Overall	Top 10 %	1	$\times 43$	3068 (21 %, 29 %)	5967 (42 %, 55 %)	1722	3565	14 322
N. Hemi.	Top 10 %	1	$\times 43$	2040 (27 %, 37 %)	2541 (34 %, 46 %)	969	1989	7539
S. Hemi.	Top 10 %	1	$\times 43$	1028 (15 %, 20 %)	3426 (51 %, 66 %)	753	1576	6783
N. Pacific	Top 14 %	1	$\times 43$	1018 (27 %, 35 %)	1343 (36 %, 47 %)	510	863	3734
N. Atlantic	Top 14 %	1	$\times 43$	891 (27 %, 38 %)	1061 (33 %, 45 %)	421	877	3250
N. Atlantic	Top 16 %	1	2011–2012	20 (24 %, 36 %)	26 (31 %, 46 %)	10	27	83
N. Atlantic	Top 28 %	2	2011–2012	18 (24 %, 36 %)	22 (30 %, 44 %)	10	24	74
N. Atlantic	All cyclones	1	2011–2012	51 (10 %, 17 %)	222 (43 %, 74 %)	29	215	517
N. Atlantic	All cyclones	2	2011–2012	33 (12 %, 20 %)	110 (42 %, 67 %)	20	102	265




Figure 4. Fraction of warm-seclusion cyclones for the full dataset and for the hemispheric and main oceanic subsets. Intensity deciles are based on maximum 850 hPa ξ_{850} (see Sect. 2.1) and are calculated separately for each hemisphere (but not for each oceanic basin).

ing cyclones. Recall that the proportions of cyclones in the top intensity decile subsets exceed 10 % in the North Atlantic and North Pacific subregions because the intensity threshold used is an absolute value calculated using cyclones in the entire Northern Hemisphere. Increasing the minimum track duration increases the proportion of cyclones found in the North Atlantic subset, implying that in general the shorter-duration cyclones are weaker. The proportions of SJP and nSJP top-decile warm-seclusion cyclones are similar for a given track constraint, again implying a strong relationship between (absolute) maximum intensity and SJP likelihood.

The SJP is also found to be present in 2011–2012 cyclones that are not diagnosed as having a warm seclusion. Indeed, for example, the proportions of SJP and nSJP cyclones for all-intensity non-warm-seclusion cyclones with a minimum track duration of 1 d are 18 % and 74 %, respectively (with the remainder being marginal cyclones), and those for the top decile subset are both 44 %. Hence, the warm-seclusion and non-warm-seclusion cyclones are similarly likely to be SJP cyclones (e.g. 17 % of all warm-seclusion cyclones have an SJP from Table 1). Conversely, similar proportions of SJP and nSJP cyclones have a warm seclusion (57 % and 58 %, respectively, for all-intensity cyclones). So, warm-seclusion cyclones are not more likely to have an SJP compared to cyclones without a warm seclusion, and SJP cyclones are not more likely to have a warm seclusion compared to nSJP cyclones. More detailed analysis of the SJP non-warm-seclusion top decile cyclones (not shown) has confirmed the existence of CSI points in the cloud head tip (as required to diagnose the SJP) in some cyclones that have been correctly diagnosed as not having a warm seclusion. To date in published studies, sting jets have only been diagnosed through detailed analysis in cyclones following the Shapiro–Keyser conceptual model evolution, with a frontal fracture and a warm seclusion at maturity. The existence of the SJP in some cyclones that do not follow this evolution suggests that, while the type of instability commonly found in sting-jet cyclones can develop, the absence of a frontal-fracture region with the associated weak stability inhibits the descent of the sting jet towards the ground and consequently inhibits its potentially damaging impact. Hence, in determining whether a cyclone may yield sting jets, structure should be considered together with cloud head instability characteristics. However, more research is needed

to needed to determine whether sting jets could occur in cyclones that do not follow the Shapiro–Keyser evolution.

When considering all minimum 2 d track duration 2011–2012 cyclones (all intensities and both with and without a warm seclusion), the percentage of SJP, nSJP, and marginal cyclones are 21 %, 68 %, and 11 %, respectively, with the sum of SJP and marginal cyclones yielding a percentage virtually identical to that of the SJP cyclones found by Hart et al. (2017) and Martinez-Alvarado et al. (2018) for the same track constraints. Hence, our results for the North Atlantic basin are consistent with those of earlier studies using the same track constraints despite the use of the newer ERA5 re-analysis and changes made to the SJP diagnostic. However, they suggest that the majority of the cyclones categorised as marginal precursor cyclones here would instead have been categorised as SJP cyclones in these earlier two-category precursor studies. For the remainder of this paper we only consider the results from the 43-year winter climatology of top intensity decile cyclones and focus on comparison between the SJP and nSJP cyclones, mainly ignoring the marginal cases to emphasise the differences between cyclones that do and do not have an SJP.

4.3 Spatial and temporal variability SJP and nSJP cyclones

The spatial distributions of SJP and nSJP cyclones are compared in Fig. 5. Considering first the track densities in the Northern Hemisphere (Fig. 5a), the location of the maximum track density of the SJP cyclones is found to the southwest of that of the nSJP cyclones in both the North Atlantic and North Pacific ocean basins. Hence, SJP cyclones travel at lower latitudes and are more frequent in the early part of the storm tracks. Despite the differences in track density structure between the SJP and nSJP cyclones, the maximum densities are fairly similar for both ocean basins. In the Southern Hemisphere the storm tracks are more zonal (Fig. 5c). The maximum track density for the SJP cyclones occurs to the north of that for the nSJP cyclones (i.e. SJP cyclones have a tendency to track at more equatorward latitudes in both hemispheres) although, unlike in the Northern Hemisphere, there is no obvious longitudinal displacement. Also unlike in the Northern Hemisphere, nSJP cyclones have substantially higher densities than SJP cyclones, with the peak value approximately double (consistent with the far fewer SJP compared to nSJP cyclones here; Table 1).

As in previous cyclone tracking studies (e.g. Hoskins and Hodges, 2002), localised “hot spots” of enhanced genesis densities are found. In the Northern Hemisphere (Fig. 5b) these occur over the Gulf Stream and Kuroshio currents for both the SJP and nSJP cyclones, although in both locations the genesis density peak for the SJP cyclones is slightly to the south of that for the nSJP cyclones, and the peak densities for the SJP cyclones are higher than those for the nSJP cyclones, especially for the Kuroshio current region. A hot

spot of enhanced genesis density is also apparent over the Rocky Mountains but only for the SJP cyclones, and additional hot spots are found for the nSJP cyclones towards the eastern ends of the storm tracks (particularly to the south and east of Greenland, so possibly linked to lee cyclogenesis). In the Southern Hemisphere (Fig. 5d) the hot spots are scattered along the Southern Hemisphere storm track, with the SJP cyclone hot spots found equatorward of those of the nSJP cyclones. Particularly notable is the enhanced genesis density over South America, which peaks more than 15° equatorward for the SJP compared to the nSJP cyclones.

The temporal variability in the cyclones is presented in Fig. 6, beginning with the monthly cyclone frequency over the 6-month extended-winter season considered (Fig. 6a). The monthly cyclone frequency has different behaviours for the SJP and nSJP cyclones. In the Northern Hemisphere (and in both the North Atlantic and North Pacific basins separately) there is a strong monthly cycle in SJP cyclones with a peak in midwinter (December and January). In contrast, there is a more uniform SJP cyclone frequency in the Southern Hemisphere. For the nSJP cyclones there is no consistent frequency variability between the Northern Hemisphere, North Atlantic, and North Pacific basins, and in the Southern Hemisphere there is a peak towards the later winter months (July and August). The inter-annual variability is shown in terms of the number of cyclones and the percentage of SJP cyclones in Fig. 6b and c, respectively. There are positive trends in cyclone numbers for all cyclone categories throughout the evaluation period, with the most robust trends being for the SJP cyclones in the North Atlantic and Northern Hemisphere and for both the SJP and nSJP cyclones in the Southern Hemisphere. Since only cyclones in the top intensity decile have been evaluated, positive trends in the number of cyclones evaluated for the SJP reflect an increase in either the total number of cyclones or the intensities of cyclones (such that more cyclones in later years are found in the top intensity decile). The percentage of cyclones that have the SJP provides an alternative metric for inter-annual cyclone variability. There are positive but not significant trends in this metric for each hemisphere and basin.

4.4 Distributions of intensity metrics of SJP and nSJP cyclones in ERA5 and observations

Further evidence for the distinctiveness of cyclones with and without the SJP is now provided by comparing intensity metrics using both the ERA5 dataset and observations. Considering first ERA5, Fig. 8 shows histograms of the following cyclone intensity metrics for the Northern and Southern Hemisphere cyclones: maximum ξ_{850} , maximum low-level (925 hPa) wind speed, minimum MSLP, a simple metric for MSLP anomaly (maximum MSLP difference within 300 km of the minimum MSLP) all at the time of maximum intensity (according to ξ_{850}), and the maximum 24 h MSLP decrease along the track. For all metrics, except for the mini-




Figure 5. Track (a, c) and genesis (b, d) densities for SJP (shading) and nSJP (black contours) cyclones for the (a, b) Northern and (c, d) Southern Hemisphere. Units are the number per unit area per season, where the unit area is equivalent to a 5° spherical cap (approximately 10^6 km^2). Shading and contours share the same intervals in all of the panels.




Figure 6. Monthly (a) and inter-annual (b, c) variations in SJP cyclone counts. Units in panel (a) are the average number of cyclones per month, in panel (b) are absolute cyclone counts, and in panel (c) are the percentage of all cyclones that are SJP cyclones. The error bars in panel (a) (for legibility, shown for SJP cyclones only) indicate 2 standard errors in the means estimated from the 43 yearly values for each month, and the legends in panels (b) and (c) indicate the linear trends of each time series, with asterisks indicating trends that are non-zero according to a two-sided Wald test at the 5% level.

imum MSLP at the time of maximum intensity in the Southern Hemisphere, there is a clear distinction between the normalised frequency distributions, with the SJP cyclones being more intense (i.e. higher values of maximum ξ_{850} , low-level wind speed, MSLP anomaly and 24 hPa decrease, and lower values of minimum MSLP). This distinction is visible in the shapes of the SJP and nSJP cyclone distributions and also in the mean values, which increase in intensity going from the set of nSJP (FALSE) cyclones to that of marginal cyclones and then to SJP (TRUE) cyclones. For the maximum ξ_{850} , minimum MSLP, and MSLP anomaly, the higher-intensity tail is also longer for SJP cyclones with this long tail particularly notable for the maximum ξ_{850} distribution in the Northern Hemisphere. Comparing the distributions for the two hemispheres, the mean values of the Southern Hemisphere cyclones for the three categories are slightly higher for maximum ξ_{850} , low-level wind speed, and 24 hPa MSLP decrease. However, the more notable difference is for minimum MSLP, with Southern Hemisphere mean values for the three categories being 10–18 hPa lower than for their Northern Hemisphere counterparts. The apparently anomalous result that there is not a clear distinction between the frequency distributions of minimum MSLP for the SJP and nSJP cyclones in the Southern Hemisphere is a consequence of the difference in their genesis and track densities (Fig. 5), with SJP cyclones typically forming closer to the Equator (where climatological MSLP is higher) and travelling more zonally for most of their life cycle. The along-track evolution of the minimum MSLP is further discussed in the following section on composite analysis (Sect. 4.5).

As the vast majority of cyclones occur over the oceans, comparison of SJP and nSJP cyclones using direct observations of near-surface wind speeds is hindered by a lack of high-resolution, high-frequency global observations. The CCMP3.0 dataset provides the best available data by blending bias-corrected ERA5 near-surface winds with scatterometer observations. To avoid including cyclones with limited or no observational input, we identify the location of the maximum CCMP3.0 wind speed within a 10° box of each cyclone centre at the time of maximum intensity and filter to just those cyclones with at least two satellite observations contributing to the blended product. Figure 7 shows that the average maximum observed wind speed at the time of maximum intensity increases from nSJP cyclones to marginal cyclones to SJP cyclones. This increase is consistent with Figs. 8b, e and 13g (for 925 hPa and 850 Pa wind speed, respectively), confirming that SJP cyclones tend to have stronger wind speeds than nSJP cyclones. Furthermore, it is found that the CCMP3.0 minus ERA5 difference in maximum near-surface wind speed is larger also for the SJP cyclones than the other two categories (not shown). This finding may be evidence that there is stronger mesoscale wind activity in the SJP cyclones, e.g. associated with sting jets, which is observed in CCMP3.0 but not resolved by ERA5. However, given the non-linear nature of the bias correction




Figure 7. Maximum CCMP3.0 wind speeds within a 10° box of each cyclone centre at the time of maximum intensity. All cyclones from the CCMP3.0 period (1993–2019) that have at least two independent scatterometer observations contributing to the blended wind speed product at the grid point of the maximum are included.

applied to the ERA5 winds to produce the CCMP3.0 background field, confirming this result requires further work.

4.5 Composite SJP and nSJP cyclone characteristics

Cyclone compositing reveals systematically distinctive characteristics of cyclones with and without an SJP. Maps of selected composite fields at chosen times relative to the time of maximum cyclone intensity and for chosen ocean basins are shown in Figs. 9–12 and S1–S5 in the Supplement, and line plots showing time series of attributes of some of these fields (e.g. composite maximum 850 hPa wind speed) are shown in Fig. 13 for all four regions and for cyclones with and without the SJP. Recall that when compass directions are used for convenience below that the cyclones have been rotated prior to compositing. While the compositing, by its nature, yields smoothed cyclone fields, the key cyclone features including the cloud head, fronts, warm sector, and deep low-pressure centre can clearly be identified in the map plots. At the time of maximum intensity, the composite SJP cyclone has notably stronger low-level winds and warmer (in θ_w) warm sector and cyclone core than the composite nSJP cyclone in all three regions (North Atlantic, North Pacific, and Southern Hemisphere) included in Fig. 9, which shows composites for the SJP cyclones in the top row and composite differences (SJP minus nSJP) in the bottom row. The MSLP centre is also deeper for the Northern Hemisphere composites but not for the Southern Hemisphere composite (consistent with Figs. 8c, f and 13a). The peak wind speed region is equatorward of the composite centre, extending from the warm sector rearwards behind the cold front towards the tip of the cloud head as well as along the cold front. A warm seclusion is evident even in the smooth SJP cyclone compos-




Figure 8. Normalised frequency plots of SJP and nSJP cyclone intensity metrics. Orange and purple bars are for SJP and nSJP cyclones, respectively, with the area covered by both sets of bars in dark red. Mean values of cyclone sets are indicated by dashed vertical lines, with the values also listed in the legend, where the TRUE, FALSE, and MARGINAL labels indicate SJP, nSJP, and marginal cyclones, respectively. Panels (a)–(c), (g), (h) and (d)–(f), (j), (k) are for the Northern and Southern Hemispheres, respectively. Intensity metrics in panels (a)–(g) and (j) are considered at the time of maximum intensity of the cyclone (according to ξ_{850}). (a, d) Track maximum spatially smoothed ξ_{850} . (b, e) Maximum wind speed at 925 hPa within a 6° geodesic great circle radius centred on the track (as a simple grid point search). (c, f) Minimum MSLP within a 5° geodesic great circle radius centred on the track using a B-spline interpolation and steepest descent minimisation. (g, j) Maximum difference in MSLP in a circle with a radius of 300 km centred at the location of minimum MSLP. (h, k) Maximum 24 h decrease in MSLP along the full length of the track; this value is set to zero in the absence of any 24 h decrease.

ite at this maximum intensity time in all the regions (although a closed θ_w contour is not evident for the North Atlantic region with the contouring used).

The differences between the composite SJP and nSJP cyclones are strongest for the North Pacific cyclones in central MSLP and peak low-level wind speed. The peak differences are found on the poleward edge, i.e. closer to the cyclone centre, of the region of strongest winds in the SJP cyclones (as

can be seen in bottom row of Fig. 9 by comparing the shaded wind differences with the stippled region indicating the location of the composite SJP cyclone peak winds) and elongate along the composite cold front. Although the wind speed differences between the composite SJP and nSJP cyclones are overwhelmingly positive, for the Southern Hemisphere cyclones there is a negative anomaly of wind speed poleward of the cyclone centre, where the cold conveyor belt travels rear-




Figure 9. Maps of the low-level composite structure of the SJP cyclones (a–c) and their differences to the nSJP cyclones (d–f) at the time of maximum intensity for the (a, d) North Atlantic, (b, e) North Pacific, and (c, f) Southern Hemisphere: (a–c) 850 hPa wind speed (shaded), MSLP (black contours) with 850 hPa θ_w (brown contours) and 700 hPa cloud (stippling indicates relative humidity with respect to ice > 80 %) and (d–f) difference fields for wind speed (shaded), MSLP (black contours), and 850 hPa θ_w (brown contours), with green stippling marking the region of winds exceeding 32 m s^{-1} in the equivalent SJP composites.

ward relative to the cyclone motion, approaching the magnitude of the positive anomaly in the peak wind region. This negative anomaly is associated with a stronger MSLP pressure gradient in this region in the composite nSJP cyclone (as can be inferred from the MSLP difference field) and so is consistent with stronger gradient-wind-balanced winds in the cold conveyor belt jet. Note, however, that this negative difference in wind speed in the Southern Hemisphere composite and also to some extent the positive differences in the peak wind speed in all three regions are also partially attributable to the slower cyclone travel speed of the composite nSJP cyclones, as discussed further below.

The evolution of the low-level North Atlantic composite cyclone structure over the 24 h prior to maximum intensity is shown in Figs. 10 and 9a, d for 24–6 and 0 h prior to the time of maximum intensity, respectively. As expected, the centre of the composite SJP cyclone deepens in MSLP as the cyclone intensifies. The warm sector (demarcated by the θ_w contour of 284 K at earlier times and 282 K at later times in the top row) narrows, mainly due to the cyclonic rotation of the cold front in these rotated composites, as the composite cyclone matures from an open wave towards a secluded cyclone. The core θ_w decreases as the cyclone matures, con-

sistent with the poleward cyclone motion (see Fig. 13d). At the start of the time period shown the composite cold front is noticeably stronger in terms of the θ_w gradient than the composite warm front. However, the warm front strengthens as the cyclone matures, such that by maturity the strength of the warm front approaches that of the cold front. The magnitude of the winds in the strong-wind region increases, and the region of peak winds moves from being located in the centre of the warm sector to straddling the cold front as the composite cyclone matures, consistent with the expected increasing dominance of the cold conveyor belt jet relative to the warm conveyor belt jet (see Fig. 1 of Hewson and Neu, 2015). The differences in central MSLP and peak wind speed between the composite SJP and nSJP cyclones increase as the cyclones approach maturity. As well as the developing central MSLP pressure difference between the composite cyclones, the MSLP is far less in the SJP compared to the nSJP composite to the northwest of the cyclone centre, with this difference largest at the earliest time shown when it exceeds 6 hPa in the domain corner (the MSLP fields can also be compared using Fig. 11, which shows the fields for the composite nSJP cyclones in the bottom row). It is hypothesised that this difference results from a greater proportion of the SJP cy-




Figure 10. Maps of the low-level composite structure of the SJP cyclones (a–d) and their differences to the nSJP cyclones (e–h) for the North Atlantic at (a, e) 24, (b, f) 18, (c, g) 12, and (d, h) 6 h prior to the time of maximum intensity. Fields are shown as in Fig. 9.




Figure 11. Maps of the composite upper-tropospheric wind structure of the SJP (a–d) and nSJP (e–h) cyclones for the North Atlantic at (a, d) 24, (b, e) 12, and (c, f) 0 h prior to the time of maximum intensity. Fields are shown as in the top row of Fig. 10 but with the 300 hPa wind field shaded instead.

clones forming as secondary cyclones, such that the lower pressure to their northeast is due to the associated primary cyclone, but this hypothesis has not been tested here. Equivalent figures to Fig. 10 are shown in Figs. S1 and S2 for the North Pacific and Southern Hemisphere basins, respectively,

and show similar evolution characteristics to the North Atlantic.

The evolution of the upper-tropospheric (300 hPa) wind structure for the composite North Atlantic SJP and nSJP cyclones is shown in Fig. 11. The upper-level jet is notably stronger in the composite SJP cyclone and while the cyclone




Figure 12. Maps of the composite 700 hPa PV structure of the SJP (a–d) and nSJP (e–h) cyclones for the North Atlantic. In panels (a)–(c) and (e)–(g), shading is the composite number of negative (symmetrically unstable) PV points at (a, e) 24, (b, f) 12, and (c, g) 0 h prior to the time of maximum intensity; in panels (d, h) shading is the composite PV at the time of maximum intensity. All other fields are as in the top row of Fig. 10.

centre moves northwards relative to the jet core as the cyclone intensifies in both sets of composites, in the SJP composite it moves from being embedded in the jet core to the left-jet exit region (a preferential region for cyclone intensification due to enhanced vortex stretching), whereas in the nSJP composite it is in the jet exit region throughout. This difference suggests that cyclones with an SJP are more likely to have crossed from the equatorward to poleward side of the jet as they intensify and so are located in the right-jet entrance region earlier in their development (another preferential region for cyclone intensification due to enhanced vortex stretching). Equivalent figures to Fig. 11 are shown in Figs. S3 and S4 for the North Pacific and Southern Hemisphere basins, respectively. The composite upper-level jet is strongest overall for SJP cyclones in the North Pacific, and the cyclone centre also begins more equatorward relative to the jet core in the SJP compared to the nSJP composites in these other basins.

The mid-level PV structures of the composite North Atlantic cyclones are compared in Fig. 12. A strong PV anomaly exists in the centre of the composites at the time of maximum intensity, extending radially towards the cloud edge to the north and east, with PV values exceeding 1 PVU (where $1 \text{ PVU} = 10^{-6} \text{ K kg}^{-1} \text{ m}^2 \text{ s}^{-1}$) (Fig. 12d and h). This localised anomaly is embedded in a general gradient of PV across the domain, from lower values in the south to higher values in the north (with extreme values towards the southwest and northeast corners of the domain). This gradient is consistent with the expected background PV gradient due

to the latitudinal dependence of the Coriolis parameter; the rotation of this background gradient from south to north is due to the rotation of the cyclones prior to compositing, as this rotation is generally slightly clockwise due to the typical northwestward cyclone motion. The anomalously large mid-level PV at the composite centres is consistent with strong production from diabatic cloud processes, a known feature of intense extratropical cyclones (e.g. Čampa and Wernli, 2012). The PV has larger values in the SJP composite, with the maximum central value (within 300 km of the composite centre) exceeding that in the nSJP composite by more than 0.2 PVU at this time (Fig. 13c), implying stronger generation from diabatic processes. There is also a difference in the background PV between the two composites, with values in the SJP composite being generally lower across the domain; the magnitude of this difference exceeds 0.5 PVU in the north of the domain. This difference arises from the generally more equatorward location (associated with smaller Coriolis parameter values) of the SJP cyclones.

The rest of the panels in Fig. 12 show the evolution of a PV measure of mesoscale instability. Although the composite PV is positive throughout the domain for both sets of composites, individual cyclones have localised regions of negative PV. Negative PV implies the existence of (dry) symmetric instability, provided that the environment is inertially and gravitationally stable, and has been previously found in individual case studies of cyclones containing sting jets (Volonté et al., 2018, 2020, 2023b; Gray and Volonté, 2024), often forming in narrow bands within the cloud head elongated




Figure 13. Time series of composite field statistics: (a) minimum MSLP, (b) maximum θ_w within 300 km of the composite MSLP centre, (c) maximum 700 hPa PV within 300 km of the composite MSLP centre (minimum value for the Southern Hemisphere), (d) storm travel direction, (e) number of cloudy points (700 hPa relative humidity with respect to ice > 80 %) within 600 km of the composite MSLP centre, (f) total number of composite mean points with symmetrically unstable 700 hPa PV (negative and positive PV for the Northern and Southern Hemisphere, respectively), (g) maximum 850 hPa wind speed, (h) 850 hPa system-relative wind speed at the location of the maximum 850 hPa (Earth-relative) wind speed, and (i) maximum 300 hPa wind speed. In panel (d) the dark and pale arrows are for the sting-jet and non-sting-jet cyclones, respectively; in all other panels the solid and dotted lines are for the sting-jet and non-sting-jet cyclones, respectively.

along the direction of the bent-back front and tracking towards the cloud head tip as the cyclone develops. Hence the existence of negative PV is an alternative, independent diagnostic for mesoscale instability to the CSI-based diagnostic that has been used to classify the cyclones according to the likely existence of a sting jet within them. The narrow and transient nature, and typically small magnitude compared to nearby regions of positive PV, of the negative PV regions means that they are not visible in the composite PV fields. Instead, here the panels show the average number of points with negative PV in the composites. For example, peak values of this metric for the North Atlantic composites (which occur in Fig. 12a) approach 0.07 (i.e. 7 %). In the SJP composites, two regions with larger values of negative PV are visible (Fig. 12a–c). The peak values are at the tip of the cloud head at 24 h prior to the time of maximum intensity and are within a band of large values that wraps around the north of the cyclone centre, aligning with the outer edge of the

cloud head. The second region of larger negative PV points lies in a band along the cold front (along the 284 K θ_w contour). There is also a local enhancement in the number of negative PV points in the region between the cloud head tip and cold front. While bands of enhanced negative PV points continue to exist aligned along the cold front at 12 and 0 h prior to the time of maximum intensity in the SJP composite, the band aligned with the cloud head decays markedly over this time period, consistent with a stronger rate of release than generation of the mesoscale instability giving rise to negative PV in this region. There are far fewer negative PV points in the cloud head and cold front regions in the nSJP of composites. Although there is some local enhancement of the nSJP composites, with values peaking within the cloud head directly north of the composite centre at 24 h prior to maximum intensity, it is clear that there is substantially more mesoscale instability in the SJP composites, as is expected given the design of the precursor diagnostic. The same com-

posite symmetrically unstable PV features and feature evolution are present to some degree in the other two basins (Fig. S5), although, relative to the North Atlantic, there are substantially more and fewer points in the SJP composites for the North Pacific and Southern Hemisphere regions, respectively. In particular, bands aligned with the cloud head and cold front exist that decay and do not decay with time, respectively, in all three basins.

Finally, time series of composite field statistics are shown in Fig. 13 for times from 36 to 0 h prior to the time of maximum composite intensity and for all four regions: the North Atlantic, North Pacific, and Northern and Southern Hemisphere. These time series enable more quantitative analysis of some previously identified characteristics (e.g. minimum MSLP evolution) and evaluation of some new diagnostics. The composite SJP cyclones deepen more rapidly than the composite nSJP cyclones in all regions from 24 h prior to maximum intensity (Fig. 13a), although, consistent with Figs. 8f and 9f, in the Southern Hemisphere the minimum MSLP pressure is similar in both composites at the time of maximum intensity as a consequence of the differences in composite MSLP at the start of the rapid deepening period. The differences in the travel direction and speed between the SJP and nSJP composites are also consistent for the four regions (Fig. 13d). The composite SJP cyclones travel faster throughout the 36 h considered and change from initially travelling in a more zonal direction, compared to the composite nSJP cyclones, to travelling in a more poleward direction as they approach their maximum intensity.

Composite SJP cyclones have a warmer core, with the maximum 850 hPa θ_w (within 300 km of the composite cyclone centre) being greater than 1.7 K (and up to 3.9 K) warmer in all regions throughout the time period shown (Fig. 13b). In all four regions, there is a much larger increase in cloudiness (measured here as the number of 700 hPa cloudy points within 600 km of the composite cyclone centre) as the composite cyclones approach maturity in those with the SJP (Fig. 13e), although, due to differences in cloudiness at the start of the time period shown, only in the Northern Hemisphere regions is the cloudiness greater in the SJP composites at the time of maximum maturity. The warmer core at the start of time period shown is consistent with the genesis regions of the SJP cyclones being displaced towards the Equator relative to those of the nSJP cyclones (Fig. 5b and d). At maximum intensity time, the SJP cyclones are found poleward of the nSJP cyclones on average (not shown), and so the warmer core is then consistent with stronger warm seclusions in the SJP cyclones. The development of increased cloudiness in the composite SJP cyclones relative to the composite nSJP cyclones in the Northern Hemisphere regions is not surprising, given that the diagnostic used to classify the precursor cyclones is based on the existence of a mesoscale instability that occurs in cloud.

Next, PV metrics are considered. The enhanced mid-level PV in the composite SJP cyclone core, relative to that

in the nSJP composite, shown for the North Atlantic at maximum intensity time in Fig. 12d and h, develops during the rapid deepening phase in all four regions (Fig. 13c, noting that of course the minimum rather than maximum PV is considered for the Southern Hemisphere). In contrast, the enhanced numbers of symmetrically unstable PV points for the composite SJP cyclones are present throughout the time period shown with the number of points slowly decreasing with time for most composites, with the possible exception of the Southern Hemisphere SJP composite (Fig. 13c). This decrease is seen despite all points within the composite being counted in Fig. 13c, but only at the cloud head tip was a strong decay seen in Figs. 12 and S5.

The bottom row of Fig. 13 shows wind speed time series. The maximum 850 hPa wind speed is clearly faster for the SJP composites, with the differences between the two composites for the same region increasing during the 36 h intensification period considered (Fig. 13g, consistent with the maps in Figs. 9, 10, S1, and S2). However, as shown in Fig. 13i (and consistent with Figs. 11, S3, and S4), the upper-tropospheric jet is also substantially faster in the precursor composite, leading to the faster travel of the SJP cyclones evident in Fig. 13d. To address the question of whether the 850 hPa winds are stronger in the SJP cyclones simply because they are travelling faster, Fig. 13h shows the 850 hPa system-relative wind speed at the same location as the maximum 850 hPa (Earth-relative) wind speed. While the system-relative wind speed increases in all the composites as the cyclones intensify, the speeds for the composites with the SJP diverge from (becoming faster than) those for the nSJP composites for each region during the last 18 h of intensification.

5 Conclusions

The aim of this study was to produce the first climatology of extratropical cyclones containing the transient mesoscale descending jet termed a sting jet and so explore their existence beyond the North Atlantic–European region, where the vast majority of previous research (including climatologies and case studies) has been focused. An algorithm for precursors to sting jets (SJPs) has been refined using a set of expertly assessed notable storms and applied to more than 10 000 tracked cyclones identified in 43 years of ERA5 data. This cyclone dataset consisted of those cyclones containing a warm seclusion, as diagnosed by a new method employing a watershed algorithm, in the top intensity decile of all tracked ERA5 cyclones. The key findings of this study address the two questions posed in Sect. 1 as follows.

1. Cyclones containing sting jets are likely to be present in all major ocean basins, despite a lack of published case analyses outside of the North Atlantic–European region.
 - Globally the percentages of the diagnosed warm-seclusion cyclones in the top intensity decile with

the SJP, without the SJP, and marginal cases are 29 %, 55 %, and 16 %, respectively. If all of the cyclones without a warm-seclusion in this top intensity decile are assumed to not yield a sting jet (because of the absence of a weak stability frontal-fracture region), then 21 % of the top intensity decile cyclones may have a sting jet. However, SJP are far more common in the Northern Hemisphere (37 % of warm-seclusion cyclones, 27 % of all cyclones) than in the Southern Hemisphere (20 %, 15 %), with similar values for the North Atlantic and North Pacific basins as for the whole Northern Hemisphere. The precise percentages will be dependent on the thresholds chosen for the SJP diagnostic (albeit having been refined using a holistically assessed set of notable storms). Analysis of all cyclones (all intensities and both with and without a diagnosed warm seclusion) for a single extended-winter season over the North Atlantic has revealed that the precursor is equally likely to be present in cyclones with and without a warm seclusion. As the sting jet is unlikely to be able to descend towards the surface in cyclones that do not follow the Shapiro–Keyser evolution with the associated frontal fracture and warm seclusion, this result implies that estimates of sting-jet prevalence based on instability metrics only (such as in Hart et al., 2017) are likely overestimates.

- SJP cyclones are more likely to have their genesis nearer the Equator than nSJP cyclones (in both hemispheres) and to form over the warm western ocean boundary currents in the Northern Hemisphere. Composite SJP cyclones track faster than nSJP cyclones, and initially track more zonally, before changing to have a greater poleward component than nSJP cyclones in all regions. In the North Atlantic, the difference in track density between SJP and nSJP cyclones found here is consistent with the previous climatology for this region (Hart et al., 2017).
 - In deriving the climatology, it was also found that the proportion of cyclones with a warm seclusion increases strongly with the decile of maximum intensity (and is similar for all regions), such that about 75 % of cyclones in the top intensity decile have warm seclusions. This result implies a strong relationship between cyclone structure and intensity.
2. Cyclones with and without the SJP have distinct characteristics in terms of their spatial and temporal variability, distributions of various intensity metrics, and composite structure. Here we reiterate that the ERA5 reanalysis data used are produced using a model that has insufficient resolution to fully represent sting jets.

However, it is possible that the assimilation of observational data, including in situ observations of 10 m winds over oceans from scatterometers, may lead to partial representation of sting-jet winds in some reanalyses, although the surface footprint of sting jets, estimated as perhaps 50–100 km across (Hewson and Neu, 2015), would barely be represented in this dataset with ~ 30 km grid spacing even if sting jets were represented. Hence, the differences found between SJP and nSJP cyclones in the ERA5 dataset are not due to sting-jet presence per se but instead reveal larger-scale differences between cyclones with and without the SJP. While specifically the SJP is a diagnosis of substantial CSI in the cyclone cloud head, more generally it likely distinguishes those cyclones with strong cloud-related diabatic processes.

- SJP cyclones are globally more intense than nSJP cyclones in terms of maximum ξ_{850} , maximum low-level wind speed, and maximum MSLP anomaly at the time of maximum intensity (in terms of ξ_{850}), as well as in terms of maximum 24 h MSLP decrease. While Northern Hemisphere SJP cyclones are on average 6 hPa deeper in MSLP at the time of maximum intensity, in the Southern Hemisphere the values are similar. The latter, apparently anomalous, result is a consequence of the different tracks typically taken by Southern Hemisphere SJP and nSJP cyclones.
- Comparison of cyclone maximum wind speeds derived using a dataset consisting of a blend of scatterometer and bias-corrected ERA5 data for available cyclones show that the median wind speed for the SJP cyclones is more than 3 m s^{-1} greater than that for the nSJP cyclones, providing independent verification of the wind speed differences between the two cyclone categories.
- Comparison of the composite structure of SJP and nSJP cyclones reveals significant structural differences. In addition to the differences highlighted above, it is particularly notable that the composite SJP cyclones have a warmer (higher values of 850 hPa θ_w) core than the nSJP cyclones in all regions and that the composite MSLP field is consistent with a greater proportion of SJP cyclones forming as secondary cyclones. The composite upper-tropospheric jet (at 300 hPa) is substantially faster for the composite SJP cyclones throughout the 36 h prior to maximum intensity time considered (typically by about 10 m s^{-1}), and SJP cyclones are more likely to cross from the equatorward to poleward side of the jet as they intensify. Analysis of the system-relative wind speeds shows that the faster (Earth-relative) low-level wind speeds found in the SJP cyclones are greater than expected due solely to the faster cyclone travel speed.

- Finally, analysis of the composite PV structure of the cyclones reveals larger maximum (minimum in Southern Hemisphere) mid-level PV values in the SJP cyclone cores, consistent with stronger PV generation through cloud latent heating. There are also more points with symmetrically unstable PV (negative in the Northern Hemisphere and positive in the Southern Hemisphere) in the SJP cyclones, providing further evidence of enhanced mesoscale instability in these cyclones. The number of points in the cloud head tip decreases with time, implying that the instability is released.

The implications of these findings are that sting-jet cyclones are ubiquitous in the extratropics (in both hemispheres) despite published detailed case studies of these cyclones to date only existing for cyclones in the North Atlantic–European regions. The potential enhancement of near-surface winds and gusts due to sting jets, which may be poorly represented in weather forecast models, has implications for weather warnings due to land-falling cyclones. The greater intensity of cyclones containing the SJP diagnostic in ERA5 data (compared to those without the SJP) and structural differences revealed by the compositing reflect the important contribution of cloud diabatic processes (leading to CSI and other mesoscale instabilities) in cyclone development, with implications for changes in the intensity of cyclones in a warmer future climate where cloud diabatic processes are likely to be enhanced.

Open questions remaining include (i) why SJP cyclones are far less prevalent in the Southern Hemisphere than in the Northern Hemisphere and the consequence of this difference for the presence of sting jets; (ii) whether sting-jet cyclones occur beyond the extratropical band considered here; (iii) the relationship between the presence (and magnitude) of the SJP, the presence of sting jet(s), and enhanced surface winds due to sting jet(s) in cyclones; and (iv) whether, despite a lack of published analysed case studies, sting jets could occur in cyclones that meet the criteria for the SJP and yet follow an evolution far removed from the classical Shapiro–Keyser evolution with a warm seclusion and frontal fracture. In particular, the latitudinal constraints used for the cyclone tracks (40–80°) exclude much of the Mediterranean Sea, where diabatic processes are known to be particularly important for the development of some cyclones (Flaounas et al., 2021) and where strongly convective medicanes with tropical characteristics occur. The differences in characteristics of the cyclones in this region compared to those in the main storm tracks, including typically smaller sizes and shorter lifetimes, mean that further work would be needed to refine and verify the SJP for this region. It is anticipated that the databases generated here of cyclones with and without SJPs will enable case studies and other analyses to be performed to address some of these questions. Finally, while outside the scope of this study, interesting questions linked to conceptual models of

cyclone development can be addressed using the generated database of intense warm-seclusion cyclones.

Code and data availability. For the ERA5 data, please see Hersbach et al. (2023a) (<https://doi.org/10.24381/cds.bd0915c6>) and Hersbach et al. (2023b) (<https://doi.org/10.24381/cds.adbb2d47>). The results contain modified Copernicus Climate Change Service information. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or the data it contains. The warm-seclusion algorithm was developed using the watershed image segmentation algorithm provided by the scikit-image Python library (van der Walt et al., 2014). The TRACK algorithm is available on the University of Reading's Git repository (GitLab) at <https://gitlab.act.reading.ac.uk/track/track> (Hodges, 2023). The satellite images used in Fig. 2 and described in Sect. 2.3 were obtained from the EUMETSAT data portal using the eumdac Python library, and the CCMP3.0 data were downloaded from Mears et al. (2022b) (<https://doi.org/10.56236/RSS-uv6h30>). Code to calculate DSCAPE from vertical profiles is available from GitHub at <https://github.com/omartineza/csisounding> (csisounding, 2019). The cyclone tracks, cyclone relative fields extracted from ERA5, derived DSCAPE diagnostics, and other derived fields are archived at the Centre for Environmental Data Analysis (CEDA; Gray et al., 2024) at <https://doi.org/10.5285/4aac4f8ba15f43e59eb81756b464c9fb>.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/wcd-5-1523-2024-supplement>.

Author contributions. SLG, OMA, and AV conceptualised the study and acquired funding for it. All authors performed parts of the software development and data analysis. SLG led the paper writing, but all authors drafted parts of the text and contributed to the editing and review process. BJH led the data curation.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. The authors thank Kevin Hodges for supplying the identification and tracks of cyclones from the ERA5 dataset as well as code to produce the cyclone density maps. We also thank Helen Dacre for her useful comments on a draft of this paper.

Financial support. This work was funded by the UKRI Natural Environment Research Council (NERC), grant number NE/X010473/1. Oscar Martínez-Alvarado and Ben J. Harvey were also funded by the NERC Climate change in the Arctic–North Atlantic region and impacts on the UK (CANARI) programme, grant number NE/W004984/1.

Review statement. This paper was edited by Lukas Papritz and reviewed by Emmanouil Flaounas and one anonymous referee.

References

- Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A cross-calibrated, multi-platform ocean surface wind velocity product for meteorological and oceanographic applications, *B. Am. Meteorol. Soc.*, 92, 157–174, 2011.
- Bjerknes, J.: On the structure of moving cyclones, *Mon. Weather. Rev.*, 47, 95–99, [https://doi.org/10.1175/1520-0493\(1919\)47<95:OTSOMC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1919)47<95:OTSOMC>2.0.CO;2), 1919.
- Brâncuș, M., Schultz, D. M., Antonescu, B., Dearden, C., and Ștefan, S.: Origin of Strong Winds in an Explosive Mediterranean Extratropical Cyclone, *Mon. Weather Rev.*, 147, 3649–3671, <https://doi.org/10.1175/MWR-D-19-0009.1>, 2019.
- Browning, K. A.: The sting at the end of the tail: Damaging winds associated with extratropical cyclones, *Q. J. Roy. Meteorol. Soc.*, 130, 375–399, <https://doi.org/10.1256/qj.02.143>, 2004.
- Browning, K. A. and Roberts, N. M.: Structure of a frontal cyclone, *Q. J. Roy. Meteorol. Soc.*, 120, 1535–1557, <https://doi.org/10.1002/qj.49712052006>, 1994.
- Browning, K. A., Smart, D. J., Clark, M. R., and Illingworth, A. J.: The role of evaporating showers in the transfer of sting-jet momentum to the surface, *Q. J. Roy. Meteorol. Soc.*, 141, 2956–2971, <https://doi.org/10.1002/qj.2581>, 2015.
- Čampa, J. and Wernli, H.: A PV perspective on the vertical structure of mature midlatitude cyclones in the Northern Hemisphere, *J. Atmos. Sci.*, 69, 725–740, <https://doi.org/10.1175/JAS-D-11-050.1>, 2012.
- Catto, J., Ackerley, D., Booth, J., Adrian J. Champion, B. A. C., Pfahl, S., Pinto, J. G., Quinting, J. F., and Seiler, C.: The Future of Midlatitude Cyclones, *Curr. Clim. Change Rep.*, 5, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>, 2019.
- Clark, P. A. and Gray, S. L.: Sting jets in extratropical cyclones: a review, *Q. J. Roy. Meteorol. Soc.*, 144, 943–969, <https://doi.org/10.1002/QJ.3267>, 2018.
- Clark, P. A., Browning, K. A., and Wang, C.: The sting at the end of the tail: Model diagnostics of fine-scale three-dimensional structure of the cloud head, *Q. J. Roy. Meteorol. Soc.*, 131, 2263–2292, <https://doi.org/10.1256/qj.04.36>, 2005.
- csisounding: Computation of DOWndraught Slantwise Convective Available Potential Energy (DSCAPE) from vertical soundings, GitHub [code], <https://github.com/omartineza/csisounding> (last access: 1 May 2024), 2019.
- Flaounas, E., Gray, S. L., and Teubler, F.: A process-based anatomy of Mediterranean cyclones: from baroclinic lows to tropical-like systems, *Weather Clim. Dynam.*, 2, 255–279, <https://doi.org/10.5194/wcd-2-255-2021>, 2021.
- Gray, S. L. and Volonté, A.: Extreme low-level wind jets in Storm Ciarán, *Weather*, 79, 384–389, <https://doi.org/10.1002/wea.7620>, 2024.
- Gray, S. L., Martínez-Alvarado, O., Baker, L. H., and Clark, P. A.: Conditional symmetric instability in sting-jet storms, *Q. J. Roy. Meteorol. Soc.*, 137, 1482–1500, <https://doi.org/10.1002/qj.859>, 2011.
- Gray, S. L., Martínez-Alvarado, O., Ackerley, D., and Suri, D.: Development of a prototype real-time sting-jet precursor tool for forecasters, *Weather*, 76, 369–373, <https://doi.org/10.1002/WEA.3889>, 2021.
- Gray, S. L., Volonté, A., Martínez-Alvarado, O., and Harvey, B.: A global climatology of sting-jet cyclones: TRACK files and Sting-Jet Precursor Cut-Outs, NERC EDS Centre for Environmental Data Analysis [data set], <https://doi.org/10.5285/4aac4f8ba15f43e59eb81756b464c9fb>, 2024.
- Hart, N. C. G., Gray, S. L., Clark, P. A., Hart, N. C. G., Gray, S. L., and Clark, P. A.: Sting-Jet Windstorms over the North Atlantic: Climatology and Contribution to Extreme Wind Risk, *J. Climate*, 30, 5455–5471, <https://doi.org/10.1175/JCLI-D-16-0791.1>, 2017.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.bd0915c6>, 2023a.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023b.
- Hewson, T. D. and Neu, U.: Cyclones, windstorms and the IMILAST project, *Tellus A*, 67, 27128, <https://doi.org/10.3402/tellusa.v67.27128>, 2015.
- Hodges, K. I.: Feature Tracking on the Unit Sphere, *Mon. Weather Rev.*, 123, 3458–3465, [https://doi.org/10.1175/1520-0493\(1995\)123<3458:FTOTUS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1995)123<3458:FTOTUS>2.0.CO;2), 1995.
- Hodges, K. I.: Spherical Nonparametric Estimators Applied to the UGAMP Model Integration for AMIP, *Mon. Weather Rev.*, 124, 2914–2932, [https://doi.org/10.1175/1520-0493\(1996\)124<2914:SNEATT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1996)124<2914:SNEATT>2.0.CO;2), 1996.
- Hodges, K. I.: Adaptive Constraints for Feature Tracking, *Mon. Weather Rev.*, 127, 1362–1373, [https://doi.org/10.1175/1520-0493\(1999\)127<1362:ACFFT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1999)127<1362:ACFFT>2.0.CO;2), 1999.
- Hodges, K. I.: TRACK tracking and analysis system for weather, climate and ocean data, Gitlab [code], <https://gitlab.act.reading.ac.uk/track/track> (last access: 13 October 2023), 2023.
- Hoskins, B. J. and Hodges, K. I.: New Perspectives on the Northern Hemisphere Winter Storm Tracks, *J. Atmos. Sci.*, 59, 1041–1061, [https://doi.org/10.1175/1520-0469\(2002\)059<1041:NPOTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPOTNH>2.0.CO;2), 2002.
- Manning, C., Kendon, E. J., Fowler, H. J., Roberts, N. M., Berthou, S., Suri, D., and Roberts, M. J.: Extreme windstorms and sting jets in convection-permitting climate simulations over Europe,

- Clim. Dynam., 58, 2387–2404, <https://doi.org/10.1007/S00382-021-06011-4>, 2021.
- Manning, C., Kendon, E. J., Fowler, H. J., and Roberts, N. M.: Projected increase in windstorm severity and contribution from sting jets over the UK and Ireland, *Weather Clim. Ext.*, 40, 100562, <https://doi.org/10.1016/j.wace.2023.100562>, 2023.
- Martínez-Alvarado, O., Gray, S. L., Catto, J. L., and Clark, P. A.: Sting jets in intense winter North-Atlantic windstorms, *Environ. Res. Lett.*, 7, 24014, <https://doi.org/10.1088/1748-9326/7/2/024014>, 2012.
- Martínez-Alvarado, O., Gray, S. L., Clark, P. A., and Baker, L. H.: Objective detection of sting jets in low-resolution datasets, *Meteorol. Appl.*, 20, 41–55, <https://doi.org/10.1002/met.297>, 2013.
- Martínez-Alvarado, O., Gray, S., Hart, N., Clark, P., Hodges, K., and Roberts, M.: Increased wind risk from sting-jet windstorms with climate change, *Environ. Res. Letts.*, 13, 044002, <https://doi.org/10.1088/1748-9326/aaae3a>, 2018.
- Mears, C., Lee, T., Ricciardulli, L., Wang, X., and Wentz, F.: Improving the accuracy of the Cross-Calibrated Multi-Platform (CCMP) ocean vector winds, *Remote Sens.*, 14, 4230, <https://doi.org/10.3390/rs14174230>, 2022a.
- Mears, C., Lee, T., Ricciardulli, L., Wang, X., and Wentz, F.: RSS Cross-Calibrated Multi-Platform (CCMP) 6-hourly ocean vector wind analysis on 0.25 deg grid, Version 3.0, Remote Sensing Systems [data set], <https://doi.org/10.56236/RSS-uv6h30>, 2022b.
- Parton, G. A., Vaughan, G., Norton, E. G., Browning, K. A., and Clark, P. A.: Wind profiler observations of a sting jet, *Q. J. Roy. Meteorol. Soc.*, 135, 663–680, <https://doi.org/10.1002/qj.398>, 2009.
- Shapiro, M. A. and Keyser, D.: Fronts, jet streams and the tropopause, in: *Extratropical cyclones: The Erik Palmén memorial volume*, edited by: Newton, C. W. and Holopainen, E. O., American Meteorological Society, 167–191, ISBN 10:999112571X, ISBN 13:978-9991125718, 1990.
- van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Gouillart, E., Yu, T., and the scikit-image contributors: scikit-image: image processing in Python, *Peer J.*, 2, e453, <https://doi.org/10.7717/peerj.453>, 2014.
- Volonté, A., Clark, P. A., and Gray, S. L.: The role of mesoscale instabilities in the sting-jet dynamics of windstorm Tini, *Q. J. Roy. Meteorol. Soc.*, 144, 877–899, <https://doi.org/10.1002/QJ.3264>, 2018.
- Volonté, A., Clark, P. A., and Gray, S. L.: Idealised simulations of cyclones with robust symmetrically unstable sting jets, *Weather Clim. Dynam.*, 1, 63–91, <https://doi.org/10.5194/wcd-1-63-2020>, 2020.
- Volonté, A., Gray, S. L., Clark, P. A., Martínez-Alvarado, O., and Ackerley, D.: Strong surface winds in Storm Eunice. Part 1: storm overview and indications of sting jet activity from observations and model data, *Weather*, 79, 40–45, <https://doi.org/10.1002/WEA.4402>, 2023a.
- Volonté, A., Gray, S. L., Clark, P. A., Martínez-Alvarado, O., and Ackerley, D.: Strong surface winds in Storm Eunice. Part 2: airstream analysis, *Weather*, 79, 54–59, <https://doi.org/10.1002/WEA.4401>, 2023b.