

Biomass partitioning and growth efficiency in four naturally regenerated forest tree species

Article

Accepted Version

Postprint - Appendix

Konopka, B., Pajtik, J., Moravcik, M. and Lukac, M. ORCID: <https://orcid.org/0000-0002-8535-6334> (2010) Biomass partitioning and growth efficiency in four naturally regenerated forest tree species. Basic and Applied Ecology, 11 (3). pp. 234-243. ISSN 1439-1791 doi: <https://doi.org/10.1016/j.baae.2010.02.004> Available at <https://centaur.reading.ac.uk/18389/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.baae.2010.02.004>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Appendix A

Table 1 Site characteristics of selected European beech plots.

Locality	ŠLP I	SLP II	Sekier I	Sekier II	Hrochoť	Kráľová	Zvolen
Altitude (m a.s.l.)	710	675	670	660	620	550	460
Longitude (°) (') (")	19°00'23"	19°03'17"	19°13'26"	19°13'18"	19°17'01"	19°09'36"	19°07'36"
Latitude (°) (') (")	48°37'56"	48°38'45"	48°29'57"	48°29'58"	48°39'30"	48°32'05"	48°33'10"
Exposition	W	SW	N	NW	NW	NW	N
Slope (%)	15	40	50	50	20	20	20
Average annual temperature (°C)	6.8	7.0	6.8	6.8	6.9	7.5	7.9
Annual precipitation (mm)	920	900	826	824	764	766	740
Soil group (FAO)	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol
Geobiocenological unit	<i>Fagetum typicum</i>	<i>Fagetum typicum</i>	<i>Fagetum typicum</i>	<i>Fagetum typicum</i>	<i>Fagetum pauper</i>	<i>Fagetum pauper</i>	<i>Querceto Fagetum</i>
Mean age (years)	9.0	6.0	9.0	4.0	8.0	6.0	3.0
Mean diameter d_0 (mm)	25.0	14.5	20.0	7.0	18.8	12.5	3.5
Mean height (m)	1.71	1.32	2.44	0.54	2.23	1.17	0.58
Canopy closure (%)	75	85	95	70	100	95	95
Basal area ($\text{m}^2 \cdot \text{ha}^{-1}$)	15.08	14.46	31.66	4.39	43.72	38.41	29.70
Number of trees in full canopy (thousand per ha)	36	92	73	146	121	251	783

Table 2 Site characteristics of selected Sessile oak plots

Locality	Antol	Ladzany I	Ladzany II	Krupina I	Krupina II	Žibritov I	Žibritov II
Altitude (m a.s.l.)	560	480	500	460	380	480	480
Longitude (°) (') (")	18°57'18"	18°51'33"	18°50'56"	19°01'21"	19°05'05"	19°01'58"	19°01'42"
Latitude (°) (') (")	48°22'12"	48°17'27"	48°17'33"	48°20'19"	48°22'44"	48°22'46"	48°22'53"
Exposition	E	SW	E	SE	N	S	S
Slope (%)	35	10	5	10	10	20	10
Average annual temperature (°C)	7.6	7.9	7.7	8.3	8.1	7.8	7.7
Annual precipitation (mm)	825	801	823	709	692	763	776
Soil group (FAO)	Cambisol	Luvisol	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol
Geobiocenological unit	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>
Mean age (years)	5.5	12.0	2.5	7.5	1.5	4.0	8.0
Mean diameter d_0 (mm)	13.6	37.9	4.1	18.6	2.7	5.9	17.6
Mean height (m)	1.01	5.07	0.26	2.00	0.16	0.47	2.99
Canopy closure (%)	90	95	75	95	60	80	95
Basal area ($\text{m}^2 \cdot \text{ha}^{-1}$)	16.21	16.95	7.20	24.40	2.65	11.43	13.64
Number of trees in full canopy (thousand per ha)	84	13	428	64	463	289	46

Table 3 Site characteristics of selected Scots pine plots

Locality	Kopčany I	Zolná	Kováčová	Kišovce	Žiar n/Hr.	Levoča	Kopčany II
Altitude (m a.s.l.)	165	430	380	640	380	620	165
Longitude (°) (') (")	17°04'55"	19°13'21"	19°04'24"	20°22'24"	18°52'33"	20°30'52"	17°04'50"
Latitude (°) (') (")	48°45'16"	48°37'11"	48°35'38"	49°01'43"	48°36'11"	49°01'07"	48°45'19"
Exposition	flat	SE	E	S	E	NE	flat
Slope (%)	0	5	15	20	15	10	0
Average annual temperature (°C)	9.9	7.7	8.0	6.2	8.2	6.2	9.9
Annual precipitation (mm)	606	714	744	639	776	644	606
Soil group (FAO)	Regosol	Luvisol	Luvisol	Leptosol	Cambisol	Cambisol	Regosol
Geobiocenological unit	<i>Carpineto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Fageto Quercetum</i>	<i>Piceeto Pinetum inferiorum</i>	<i>Fageto Quercetum</i>	<i>Pinetum dealpinum superiorum</i>	<i>Carpineto Quercetum</i>
Mean age (years)	3.0	5.0	5.0	6.0	6.0	8.0	8.0
Mean diameter d_0 (mm)	14.9	28.6	37.4	21.2	43.6	41.3	36.9
Mean height (m)	0.68	1.33	1.50	1.31	2.20	2.63	3.02
Canopy closure (%)	82	92	75	77	85	90	100
Basal area ($m^2 \cdot ha^{-1}$)	12.56	23.99	18.23	14.77	26.47	31.14	55.66
Number of trees in full canopy (thousand per ha)	70	31	16	36	18	19	43

Table 4 Site characteristics of selected Norway spruce plots

Locality	Poľana I	Poľana II	Drakšiar I	Drakšiar II	Bacúch	Č. Váh I	Č. Váh II
Altitude (m a.s.l.)	985	790	625	635	840	820	830
Longitude (°) (') (")	19°31'05"	19°32'15"	19°47'29"	19°47'37"	19°49'00"	19°57'52"	19°57'54"
Latitude (°) (') (")	48°38'31"	48°38'32"	48°50'58"	48°50'59"	48°54'05"	48°59'04"	48°59'03"
Exposition	E	SW	SW	SW	S	NE	NE
Average annual temperature (°C)	5.3	6.0	6.3	6.3	4.8	5.0	5.0
Annual precipitation (mm)	982	902	865	867	1021	865	865
Slope (%)	20	10	5	5	45	60	20
Soil group (FAO)	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol	Cambisol	Podsol
Geobiocenological unit	<i>Fagetum typicum</i>	<i>Fagetum typicum</i>	<i>Abieto Fagetum inferiorum</i>	<i>Abieto Fagetum inferiorum</i>	<i>Abieto Fagetum superiorum</i>	<i>Fagetum abietino-piceosum</i>	<i>Fagetum abietino-piceosum</i>
Mean age (years)	9.0	4.0	4.5	7.0	1.5	9.5	8.5
Mean diameter d_0 (mm)	25.7	5.2	11.6	15.6	4.6	24.5	17.2
Mean height (m)	2.13	0.34	0.68	0.92	0.22	1.81	1.29
Canopy closure (%)	95	80	80	70	30	90	95
Basal area ($m^2 \cdot ha^{-1}$)	28.5	4.7	8.8	10.7	1.5	28.4	34.2
Number of trees in full canopy (thousand per ha)	58	279	104	80	293	67	155

Table 5 European beech; regression coefficients b_0 , b_1 , b_2 , their standard errors (S.E.), p-value (P), coefficient of determination (R^2), mean squared error (MSE), logarithmic transformation bias λ and its standard deviation (S.D.) for Eqs. (1)-(3)

Eq.	Tree compartment	b_0	S.E.	P	b_1	S.E.	P	b_2	S.E.	P	R^2	MSE	λ	S.D.
(1)	Stem	-4.034	0.164	<0.001	2.852	0.063	<0.001				0.923	0.167	1.0836	0.438
	Branches	-5.982	0.218	<0.001	3.117	0.084	<0.001				0.891	0.295	1.142	0.592
	Leaves	-3.750	0.183	<0.001	2.375	0.071	<0.001				0.871	0.207	1.102	0.486
	Roots	-2.960	0.179	<0.001	2.361	0.069	<0.001				0.874	0.199	1.098	0.495
	Aboveground	-3.288	0.144	<0.001	2.777	0.056	<0.001				0.937	0.129	1.063	0.364
	Whole tree	-2.521	0.132	<0.001	2.639	0.051	<0.001				0.941	0.108	1.053	0.336
(2)	Stem	3.108	0.043	<0.001	2.302	0.073	<0.001				0.854	0.318	1.166	0.672
	Branches	1.833	0.069	<0.001	2.324	0.117	<0.001				0.703	0.803	1.474	1.671
	Leaves	2.206	0.058	<0.001	1.712	0.098	<0.001				0.643	0.573	1.308	1.068
	Roots	2.964	0.061	<0.001	1.651	0.102	<0.001				0.607	0.620	1.298	0.949
	Aboveground	3.669	0.048	<0.001	2.161	0.082	<0.001				0.806	0.396	1.216	0.855
	Whole tree	4.094	0.050	<0.001	2.000	0.085	<0.001				0.767	0.427	1.226	0.853
(3)	Stem	-1.530	0.163	<0.001	1.848	0.065	<0.001	1.015	0.054	<0.001	0.975	0.054	1.026	0.222
	Branches	-4.768	0.364	<0.001	2.630	0.144	<0.001	0.423	0.121	<0.001	0.901	0.270	1.130	0.560
	Leaves	-3.286	0.317	<0.001	2.188	0.126	<0.001	0.188	0.105	0.076	0.873	0.205	1.100	0.479
	Roots	-2.898	0.314	<0.001	2.336	0.124	<0.001	0.025	0.104	0.809	0.874	0.200	1.098	0.493
	Aboveground	-1.486	0.186	<0.001	2.054	0.074	<0.001	0.731	0.062	<0.001	0.966	0.070	1.034	0.266
	Whole tree	-1.236	0.197	<0.001	2.124	0.078	<0.001	0.521	0.065	<0.001	0.957	0.079	1.038	0.285

Table 6 Sessile oak; regression coefficients b_0 , b_1 , b_2 , their standard errors (S.E.), p-value (P), coefficient of determination (R^2), mean squared error (MSE), logarithmic transformation bias λ and its standard deviation (S.D.) for Eqs. (1)-(3)

Eq.	Tree compartment	b_0	S.E.	P	b_1	S.E.	P	b_2	S.E.	P	R^2	MSE	λ	S.D.
(1)	Stem	-4.311	0.224	<0.001	2.959	0.071	<0.001				0.942	0.240	1.121	0.544
	Branches	-6.025	0.215	<0.001	2.963	0.068	<0.001				0.947	0.221	1.107	0.496
	Leaves	-5.954	0.221	<0.001	2.768	0.070	<0.001				0.936	0.234	1.106	0.459
	Roots	-1.954	0.150	<0.001	2.066	0.046	<0.001				0.941	0.124	1.065	0.401
	Aboveground	-3.952	0.206	<0.001	2.931	0.065	<0.001				0.950	0.203	1.099	0.469
	Whole tree	-2.612	0.175	<0.001	2.646	0.005	<0.001				0.956	0.146	1.070	0.382
(2)	Stem	3.471	0.045	<0.001	2.516	0.046	<0.001				0.965	0.147	1.074	0.419
	Branches	1.839	0.086	<0.001	2.388	0.089	<0.001				0.871	0.535	1.271	0.896
	Leaves	1.370	0.074	<0.001	2.269	0.077	<0.001				0.891	0.398	1.190	0.700
	Roots	3.511	0.053	<0.001	1.719	0.051	<0.001				0.898	0.213	1.112	0.553
	Aboveground	3.769	0.050	<0.001	2.468	0.052	<0.001				0.954	0.186	1.095	0.483
	Whole tree	4.369	0.051	<0.001	2.207	0.053	<0.001				0.942	0.189	1.099	0.503
(3)	Stem	-0.066	0.170	0.698	1.324	0.063	<0.001	1.491	0.053	<0.001	0.993	0.029	1.014	0.171
	Branches	-4.263	0.432	<0.001	2.284	0.161	<0.001	0.619	0.135	<0.001	0.956	0.186	1.092	0.473
	Leaves	-3.577	0.412	<0.001	1.852	0.153	<0.001	0.835	0.129	<0.001	0.954	0.169	1.077	0.399
	Roots	-0.226	0.289	0.436	1.398	0.107	<0.001	0.611	0.091	<0.001	0.956	0.092	1.047	0.333
	Aboveground	-0.201	0.194	0.303	1.486	0.072	<0.001	1.317	0.061	<0.001	0.991	0.038	1.019	0.199
	Whole tree	0.389	0.201	0.056	1.490	0.075	<0.001	1.054	0.063	<0.001	0.989	0.040	1.020	0.209

Table 7 Scots pine; regression coefficients b_0 , b_1 , b_2 , their standard errors (S.E.), p -value (P), coefficient of determination (R^2), mean squared error (MSE), logarithmic transformation bias λ and its standard deviation (S.D.) for Eqs. (1)-(3)

Eq.	Tree compartment	b_0	S.E.	P	b_1	S.E.	P	b_2	S.E.	P	R^2	MSE	λ	S.D.
(1)	Stem	-3.143	0.122	<0.001	2.480	0.038	<0.001				0.960	0.131	1.067	0.395
	Branches	-3.985	0.125	<0.001	2.515	0.039	<0.001				0.960	0.138	1.072	0.435
	Needles	-1.961	0.144	<0.001	2.014	0.045	<0.001				0.920	0.183	1.089	0.456
	Roots	-3.628	0.130	<0.001	2.236	0.041	<0.001				0.946	0.149	1.079	0.479
	Aboveground	-1.698	0.097	<0.001	2.290	0.030	<0.001				0.970	0.083	1.041	0.306
	Whole tree	-1.531	0.089	<0.001	2.278	0.028	<0.001				0.975	0.069	1.035	0.282
(2)	Stem	3.958	0.043	<0.001	2.389	0.057	<0.001				0.912	0.293	1.156	0.648
	Branches	3.246	0.062	<0.001	2.304	0.081	<0.001				0.824	0.601	1.323	1.003
	Needles	3.843	0.060	<0.001	1.805	0.078	<0.001				0.755	0.560	1.273	0.852
	Roots	2.824	0.067	<0.001	1.965	0.087	<0.001				0.745	0.694	1.387	1.173
	Aboveground	4.880	0.050	<0.001	2.134	0.065	<0.001				0.862	0.388	1.200	0.725
	Whole tree	5.013	0.051	<0.001	2.108	0.066	<0.001				0.854	0.402	1.209	0.752
(3)	Stem	-0.702	0.152	<0.001	1.618	0.052	<0.001	0.944	0.052	<0.001	0.987	0.045	1.023	0.231
	Branches	-3.184	0.257	<0.001	2.232	0.089	<0.001	0.310	0.088	<0.001	0.962	0.129	1.066	0.403
	Needles	-1.901	0.307	<0.001	1.993	0.106	<0.001	0.024	0.105	0.822	0.920	0.184	1.089	0.457
	Roots	-4.105	0.273	<0.001	2.405	0.094	<0.001	-0.184	0.093	0.050	0.947	0.146	1.078	0.487
	Aboveground	-0.466	0.177	0.009	1.856	0.061	<0.001	0.476	0.060	<0.001	0.978	0.061	1.031	0.269
	Whole tree	-0.502	0.167	0.003	1.914	0.057	<0.001	0.398	0.057	<0.001	0.980	0.054	1.028	0.253

Table 8 Norway spruce; regression coefficients b_0 , b_1 , b_2 , their standard errors (S.E.), p -value (P), coefficient of determination (R^2), mean squared error (MSE), logarithmic transformation bias λ and its standard deviation (S.D.) for Eqs. (1)-(3)

Eq.	Tree compartment	b_0	S.E.	P	b_1	S.E.	P	b_2	S.E.	P	R^2	MSE	λ	S.D.
(1)	Stem	-3.084	0.069	<0.001	2.459	0.024	<0.001				0.986	0.090	1.046	0.324
	Branches	-3.452	0.099	<0.001	2.482	0.035	<0.001				0.971	0.185	1.093	0.488
	Needles	-2.589	0.099	<0.001	2.317	0.035	<0.001				0.967	0.185	1.082	0.397
	Roots	-2.460	0.070	<0.001	2.114	0.025	<0.001				0.980	0.095	1.046	0.309
	Aboveground	-1.842	0.069	<0.001	2.398	0.024	<0.001				0.985	0.088	1.043	0.296
	Whole tree	-1.432	0.060	<0.001	2.333	0.021	<0.001				0.988	0.067	1.033	0.258
(2)	Stem	4.010	0.035	<0.001	2.398	0.033	<0.001				0.972	0.178	1.092	0.484
	Branches	3.704	0.056	<0.001	2.376	0.054	<0.001				0.930	0.453	1.245	0.921
	Needles	4.090	0.058	<0.001	2.200	0.055	<0.001				0.914	0.486	1.232	0.816
	Roots	3.640	0.052	<0.001	2.004	0.049	<0.001				0.919	0.386	1.198	0.748
	Aboveground	5.072	0.047	<0.001	2.307	0.044	<0.001				0.949	0.308	1.160	0.684
	Whole tree	5.304	0.047	<0.001	2.239	0.044	<0.001				0.948	0.300	1.157	0.676
(3)	Stem	-0.469	0.201		1.555	0.070	<0.001	0.913	0.068	<0.001	0.994	0.041	1.020	0.205
	Branches	-2.553	0.417	<0.001	2.171	0.144	<0.001	0.313	0.141	0.028	0.972	0.180	1.089	0.460
	Needles	-2.487	0.426	<0.001	2.282	0.147	<0.001	0.036	0.144	0.805	0.967	0.186	1.082	0.396
	Roots	-2.869	0.304	<0.001	2.254	0.105	<0.001	-0.142	0.103	0.169	0.980	0.094	1.045	0.306
	Aboveground	-0.696	0.281	0.014	2.002	0.097	<0.001	0.400	0.095	<0.001	0.987	0.079	1.037	0.274
	Whole tree	-0.579	0.251	0.023	2.039	0.087	<0.001	0.297	0.085	<0.001	0.989	0.062	1.030	0.248

Fig. 1 Biomass allocation coefficients based on tree height (m) at three levels of height increment of 0.1, 0.3 and 0.5 m per annum.

Fig. 2 Inter-specific comparison of growth efficiency of stem (GE I, g g^{-1}) and woody biomass (GE II, g g^{-1}) at medium range height increment (0.3 m y^{-1}).

Fig. 3 Growth efficiency of stem (GE I, g g^{-1}) and woody biomass (GE II, g g^{-1} , left axis) at three levels height (h, m) increment and dry foliage biomass (DW, g, right axis).

Fig. 1




Fig. 2




Fig. 3

