

Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Lauer, A., Eyring, V., Righi, M., Buchwitz, M., Defourny, P., Evaldsson, M., Friedlingstein, P., de Jeu, R., de Leeuw, G., Loew, A., Merchant, C. J. ORCID: https://orcid.org/0000-0003-4687-9850, Müller, B., Popp, T., Reuter, M., Sandven, S., Senftleben, D., Stengel, M., van Roozendael, M., Wenzel, S. and Willen, U. (2017) Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool. Remote Sensing of Environment, 203. pp. 9-39. ISSN 0034-4257 doi: https://doi.org/10.1016/j.rse.2017.01.007 Available at https://centaur.reading.ac.uk/68615/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

To link to this article DOI: http://dx.doi.org/10.1016/j.rse.2017.01.007

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the <u>End User Agreement</u>.



www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

- Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data 1 using the ESMValTool 2 3 Axel Lauer^{a,*}, Veronika Eyring^a, Mattia Righi^a, Michael Buchwitz^b, Pierre Defourny^c, Martin 4 Evaldsson^d, Pierre Friedlingstein^e, Richard de Jeu^f, Gerrit de Leeuw^{g,h}, Alexander Loewⁱ, 5 6 Christopher J. Merchant^j, Benjamin Müllerⁱ, Thomas Popp^k, Maximilian Reuter^b, Stein Sandven¹, 7 Daniel Senftleben^a, Martin Stengel^m, Michel Van Roozendaelⁿ, Sabrina Wenzel^a, and Ulrika 8 Willén^d 9 ^a Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, 10 Oberpfaffenhofen, Germany 11 ^b Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany 12 13 ^c Université catholique de Louvain, Earth and Life Institute, Belgium ^d Swedish Meteorological and Hydrological Institute (SMHI), 60176 Norrköping, Sweden 14 ^e College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, 15 16 EX4 4QE, UK ^f Transmissivity/VanderSat, Space Technology Center, Noordwijk, Netherlands 17 18 ^g Finnish Meteorological Institute (FMI), Climate Research Unit, 00101 Helsinki, Finland ^h Department of Physics, University of Helsinki, 00014 Helsinki, Finland 19 Department of Geography, Ludwig Maximilians Universität, Munich, Germany 20 i 21 ^j University of Reading, Reading, UK ^k Deutsches Zentrum für Luft- und Raumfahrt (DLR), Deutsches Fernerkundungsdatenzentrum, 22 23 Oberpfaffenhofen, Germany Nansen Environmental and Remote Sensing Center, Bergen, Norway 24 1
- 25 ^m Deutscher Wetterdienst (DWD), Offenbach, Germany
- 26 ⁿ Belgian Institute for Space Aeronomy, Brussels, Belgium

^{*} Corresponding author email: axel.lauer@dlr.de

27 Abstract

28 The Coupled Model Intercomparison Project (CMIP) is now moving into its sixth phase and aims at a more routine evaluation of the models as soon as the model output is published to the 29 30 Earth System Grid Federation (ESGF). To meet this goal the Earth System Model Evaluation 31 Tool (ESMValTool), a community diagnostics and performance metrics tool for the systematic 32 evaluation of Earth system models (ESMs) in CMIP, has been developed and a first version (1.0) 33 released as open source software in 2015. Here, an enhanced version of the ESMValTool is 34 presented that exploits a subset of Essential Climate Variables (ECVs) from the European Space 35 Agency's Climate Change Initiative (ESA CCI) Phase 2 and this version is used to demonstrate 36 the value of the data for model evaluation. This subset includes consistent, long-term time series 37 of ECVs obtained from harmonized, reprocessed products from different satellite instruments for 38 sea surface temperature, sea ice, cloud, soil moisture, land cover, aerosol, ozone, and greenhouse 39 gases. The ESA CCI data allow extending the calculation of performance metrics as summary 40 statistics for some variables and add an important alternative data set in other cases where 41 observations are already available. The provision of uncertainty estimates on a per grid basis for 42 the ESA CCI data sets is used in a new extended version of the Taylor diagram and provides 43 important additional information for a more objective evaluation of the models. In our analysis 44 we place a specific focus on the comparability of model and satellite data both in time and space. 45 The ESA CCI data are well suited for an evaluation of results from global climate models across 46 ESM compartments as well as an analysis of long-term trends, variability and change in the 47 context of a changing climate. The enhanced version of the ESMValTool is released as open 48 source software and ready to support routine model evaluation in CMIP6 and at individual 49 modeling centers.

50 1 Introduction

51 Earth system models (ESMs) are essential tools for improving our understanding of the climate 52 system as well as for assessing the response of the climate system to different natural or 53 anthropogenic perturbations. Understanding of the capabilities and limitations of ESMs is a 54 cornerstone for the interpretation of model results as well as for improving the models and is obtained through a comprehensive evaluation of the models with observations. Both improved 55 56 models and an improved process understanding of the climate are important steps towards 57 reducing the uncertainties in projections of future climate change and providing more 58 trustworthy information for policy guidance. The number of models participating in the Coupled 59 Model Intercomparison Project (CMIP) that is internationally coordinating ESM simulations is 60 growing and the models participating are increasing in complexity and resolution. Traceable 61 evaluation of the CMIP model ensemble with observations is therefore a challenging task.

The experimental design of the sixth phase of the Coupled Model Intercomparison Project (CMIP6) is now finalized. A central goal of CMIP6 is an improved and more routine evaluation of the participating climate models with observations (Eyring et al., 2016a). The CMIP Diagnostic, Evaluation and Characterization of Klima (DECK) experiments and CMIP historical simulations will provide the basis for the documentation of the model simulation characteristics. The aim is in particular to diagnose and improve the understanding of the origins and consequences of systematic model errors and inter-model spread.

To support this goal, the Earth System Model Evaluation Tool (ESMValTool, Eyring et al., 2016b) has been developed. The ESMValTool is a community diagnostics and performance metrics tool for systematic evaluation of Earth system models in CMIP, which has been developed by multiple institutions in several international projects. A first version of the 73 ESMValTool has been released as open source software in 2015 and is rapidly developing to 74 include additional evaluation diagnostics and technical improvements. The ESMValTool will be 75 - together with other software packages such as the Program for Climate Model Diagnostics and 76 Intercomparison (PCMDI) metrics package (PMP, Gleckler et al., 2016) and the NCAR Climate 77 Variability Diagnostic Package (CVDP, Phillips et al., 2014) that is included in the ESMValTool 78 as a separate namelist - applied to CMIP6 results to provide a broad and comprehensive characterization of the CMIP6 models as soon as the output is published to the Earth System 79 80 Grid Federation (ESGF). The foundation that will enable this is an efficient infrastructure 81 (Evring et al., 2016c) and the community-based experimental protocols and conventions of 82 CMIP, including their extension to obs4MIPs (Teixeira et al., 2014; Ferraro et al., 2015) and 83 ana4MIPs (https://www.earthsystemcog.org/projects/ana4mips/).

84 The Climate Change Initiative of the European Space Agency (ESA CCI) is a large international 85 effort that provides global, long-term satellite data sets to the climate community that can be 86 used to evaluate and improve the models (Hollmann et al., 2013). The ESA CCI is exploiting a 87 large number of satellite observations to create robust long-term global records of selected 88 essential climate variables (ECVs; GCOS, 2010; Bojinski et al., 2014) from numerous satellites 89 and instruments. In this study, a subset of the ESA CCI Phase 2 ECVs has been implemented 90 into the ESMValTool. This enhanced version of the ESMValTool is then used to evaluate 91 CMIP5 models. ESA CCI data sets implemented so far include sea surface temperature, sea ice, 92 cloud, soil moisture, land cover, aerosol, ozone, and greenhouse gases.

93 This paper is organized as follows: section 2 provides a brief description of the ESA CCI data 94 used in this study to evaluate CMIP5 models. Section 3 summarizes the models and model 95 simulations that are evaluated with the ESA CCI and other data, section 4 demonstrates the usage

96 of the implemented ESA CCI data in summary statistics applied to CMIP5 models by calculating 97 relative space-time root-mean-square deviations (RMSDs) from climatological mean seasonal 98 cycles of selected ECVs. A specific focus is placed on the consideration of uncertainty 99 information provided with the ESA CCI data, which is displayed in extended Taylor diagrams 100 (Taylor, 2001) that are widely used to assess the performance of large model ensembles in 101 reproducing observed quantities. Further insights into the evaluation of CMIP5 models with ESA 102 CCI data and comparisons of ESA CCI data with alternative observational data sets are presented 103 in section 5. A summary and a discussion of the main results and conclusions are given in section 104 6.

105 **2** Brief description of the ESA CCI data

The datasets from the ESA CCI Phase 2 implemented into the enhanced version of the ESMValTool presented in this study are briefly described in the following. We would like to note that these datasets are only a subset of all CCIs available. It is planned to implement additional CCIs such as ocean color, sea level, ice sheets and fire as well as additional ECVs from the CCIs included here into future releases of the ESMValTool.

111 **2.1** Sea surface temperature

The ESA CCI sea surface temperature (SST) data set (Merchant et al., 2014a,b) provides multidecadal products of SST derived from infrared brightness temperatures measured from satellites. SST products (Rayner et al., 2015) are generated at full sensor resolution (1 to > 4 km) and are averaged on a regular latitude-longitude grid (0.05°). A gap-filled (Level 4 SST analysis) product covering the time 1992-2010 is currently used with the ESMValTool diagnostics. The Level 4 (L4) SST analysis is a daily 3-dimensional variational analysis of satellite data with a grid resolution of 0.05°. The analysis system is the Operational Sea surface Temperature and sea-Ice

119 Analysis (OSTIA) with improved covariance parameterization (Roberts-Jones et al., 2016). The 120 L4 SST analysis has relatively good feature resolution, which is nonetheless lower than the grid 121 resolution, and varies with the density of satellite coverage (Reynolds et al., 2013). Unlike the 122 operational OSTIA products (Donlon et al., 2012) and the older OSTIA-based observational re-123 processing (Roberts-Jones et al., 2012), no in situ data are used in this CCI product. The product 124 represents the daily value of SST at a nominal depth of 20 cm, representative of the SST 125 measured by drifting buoys and bucket observations. This is possible because the lower-level 126 SST CCI products contain both the skin (radiometric) temperature of the ocean surface at the 127 time of satellite observation estimated based on radiative transfer physics (e.g., Embury et al., 128 2012a), and a turbulence-model-based adjustment to the 20 cm depth SST at a standardized time 129 of day. The adjusted SST estimate is used as input to the L4 SST analysis. This means that the 130 L4 SST analysis can be treated as independent of in situ data, and useful as a comparison point 131 for the many SST products that are tuned to and/or incorporate in situ data. The standardization 132 of the adjustment with respect to time of day is intended to reduce aliasing of the diurnal cycle 133 into false long-term trends, as satellite overpass times vary (Embury et al., 2012b). All SSTs are 134 provided with estimates of total uncertainty, and the L4 SST analysis product includes an 135 operationally produced estimate of sea ice concentration (Good and Rayner, 2014).

Merchant et al. (2014a,b) provide an assessment of the accuracy of this product by comparison with more than 2.4 million buoys from different observational networks. A global median difference against drifting buoys of +0.05 K is observed, with a standard deviation (including the ~ 0.2 K uncertainty in the drifting buoy measurements) of 0.28 K. The comparison with Argo measurements at ~ 5 m depth (only from the latter part of the record) gives +0.04 K and 0.26 K respectively. Systematic regional errors on spatial scales of ~ 1000 km range from -0.5 K to +0.5 K, with positive bias of +0.09 K across equatorial regions overall (relative to measurements of
the global tropical moored buoy array). Regions persistently affected by mineral atmospheric
aerosol, particularly Saharan dust, appear negatively biased.

145 **2.2** Sea ice

146 The ESA CCI sea ice data set provides observational data for sea ice concentration (sic) and sea 147 ice thickness (sit) that are based on satellite retrievals for both Arctic and Antarctic sea ice. The 148 sic data set is based on passive microwave data from Special Sensor Microwave Imager (SSM/I) 149 covering the time period 1992 to 2008 and the Advanced Microwave Scanning Radiometer -150 Earth Observing System (AMSR-E) covering the time period 2003-2010 (Lavergne and Rinne, 151 2014). The data sets are provided as daily gridded sic fields for both northern hemisphere and 152 southern hemisphere on an equal area grid with 25 km grid spacing. Separate data sets for SSM/I 153 and AMSR-E are available, where the SSM/I product is more mature, while the AMSR-E data 154 can provide higher-resolution products. In addition, daily maps of total standard error and quality 155 control flags are provided. The ESA CCI sea ice data set is built upon the algorithms and 156 processing software originally developed at the EUMETSAT OSI SAF for their sic data set (RD-157 11). The algorithm used to produce the sic data sets is based on an extensive algorithm 158 intercomparison study (Ivanova et al., 2015), aiming at identifying the optimal algorithm for 159 producing sic data sets. In their study, a systematic comparison of 30 algorithms was done for 160 different ice conditions, seasons and regions. The result was an implementation of a new 161 algorithm for sic retrieval. It is based on a combination of previous algorithms and use of 162 dynamic tie points and atmospheric correction of input brightness temperatures. Error sources of 163 the sic products are particularly related to the marginal ice zone, areas of thin ice, melt-ponds in 164 the summer season (Kern et al., 2016) and land contamination in coastal regions.

165 So far, only sea ice concentration and its standard error from the ESA CCI sea ice data set have 166 been implemented into the ESMValTool. Sea ice thickness data sets from radar altimeter are also 167 developed in the CCI project, but a final data set is not yet available. The ice thickness retrieval 168 is based on sea ice freeboard measurements from altimeter that are converted to thickness using 169 the hydrostatic equilibrium assumption and a priori knowledge about snow thickness, snow and 170 ice density and penetration depth of the radar signal (Kern et al., 2015). The first ice thickness 171 data set from ENVISAT for the period 2002 to 2012 has been presented (Lavergne and Rinne, 172 2014) as monthly mean thickness for the winter months in the Arctic. There are significant 173 uncertainties in these results so far, which requires further studies to obtain a reliable product. 174 Results from CryoSat thickness retrievals from 2010 to present, however, show promising results 175 (e.g., Ricker et al., 2014, Kwok and Cunningham, 2015).

176 **2.3 Cloud**

177 The ESA CCI cloud data sets contain cloud property data retrieved from the passive satellite 178 imager sensors AVHRR, MODIS, ATSR-2, AATSR and MERIS (Stengel et al., 2016a). 179 Depending on the particular data set, time periods of 9 to 33 years between 1982 and 2014 are 180 covered. In this study we used the Cloud cci AVHRR-PM v2.0 data set (Stengel et al., 2016b), 181 which is composed of data from AVHRR on-board NOAA-7, -9, -11, -14, -16, -18 and -19 and 182 represents a nearly seamless time series from 1982 through 2014. The ESA CCI cloud data sets 183 include cloud fraction (or cloud mask), thermodynamic phase, cloud top pressure (also converted 184 to temperature and height), cloud optical thickness, cloud effective radius, cloud albedo and 185 cloud liquid/ice water path. Various processing levels are available from Level 2 (pixel-based 186 data) to daily sampled data (Level 3U) and monthly averages and histograms (Level 3C). All 187 cloud properties are accompanied by pixel-based uncertainty estimates. While for most variables these estimates are based on optimal estimation theory, cloud mask uncertainty is based on hit rate scores against measurements from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP). All pixel level uncertainties are propagated in a mathematically consistent way into the Level 3C products.

In this study monthly mean cloud fraction data (inferred from Level 3C data product with 0.5° resolution on a latitude-longitude grid) are used for comparison with CMIP5 model results. Cloud fraction represents the monthly summary of the results of Community Cloud retrieval for CLimate (CC4CL) cloud detection scheme (Sus et al., 2016; McGarragh et al., 2016). The monthly mean cloud detection uncertainty is also inferred from Level 3C products.

197 CC4CL cloud detection results have been validated against CALIOP space-based lidar
198 measurements, with a global Kuipers score of 0.66 and a global hit rate of 81% (Karl-Göran
199 Karlsson, personal communication) demonstrating the high quality of the cloud detection in the
200 AVHRR-PM v2.0 data set.

201 It needs to be noted, that the Cloud cci AVHRR-PM data set has a few limitations of which 202 particularly the underrepresentation of optically very thin clouds (with optical thicknesses of 203 below 0.15) and the sparse temporal sampling (twice a day for non-polar regions) is of relevance 204 when using this data set for model evaluation. Particularly difficult conditions for cloud detection 205 are polar night periods, for which the detection scores decrease significantly in the current 206 version of the data set. Furthermore, the monthly cloud fraction data and the corresponding 207 uncertainties of the Cloud cci AVHRR-PM data set used in this study have not undergone any 208 further processing such as satellite drift correction.

209 2.4 Soil moisture

210 The ESA CCI soil moisture product is the first ever multi-decadal satellite-based soil moisture 211 product and is currently available for the time period 1978-2015 on a daily basis and at a spatial 212 resolution of 0.25°x0.25°. The ESA CCI product represents soil moisture of the first centimeters 213 of the soil and has been generated by merging active and passive microwave-based soil moisture 214

products from multiple satellite missions (Liu et al., 2011, 2012; Wagner et al., 2012).

215 Dorigo et al. (2014) provide a comprehensive validation of the ESA CCI soil moisture using 932 216 in situ observation sites from 29 different observing networks (Dorigo et al., 2011, 2013). 217 Despite the large difficulties in validating coarse resolution satellite soil moisture products with 218 in situ point like observations (Crow et al., 2012), they conclude that the ESA CCI soil moisture product has an average unbiased root-mean-square error (RMSE) of 0.05 m³ m⁻³. It was shown 219 220 that trends in the CCI observations largely agree with those obtained from various reanalysis 221 products as well as precipitation, and vegetation vigor observations (Albergel et al., 2012; 222 Dorigo et al., 2012). In addition, over the last seven years the ESA CCI soil moisture data set has 223 been used for the yearly State of the Climate Reports issued by the National Oceanic and 224 Atmospheric Administration (NOAA; e.g., Dorigo et al., 2016). Within these studies strong 225 similarities were found between the spatial annual anomalies of ESA CCI soil moisture, and the 226 terrestrial water storage from the Gravity Recovery and Climate Experiment (GRACE; e.g., 227 Willet et al., 2014).

228 The ESA CCI soil moisture data set provides a multitude of quality flags and only soil moisture 229 estimates considered reliable are used to create the data product. Snow covered areas and frozen 230 ground are typically masked as well as dense or heterogeneously vegetated areas with high optical depth that are not expected to provide reliable soil moisture estimates (Loew, 2008;
Parinussa et al., 2011).

233 **2.5 Land cover**

234 The ESA CCI land cover time series is the first consistent 300 m global land cover data set 235 providing a characterization of the land surface from 1998 to 2012. The ESA CCI land cover 236 product (v1.6.1) corresponds to high resolution global land cover information representative of 237 three 5-year periods, referred to as epochs, for 2000 (1998-2002), 2005 (2003-2007) and 2010 238 (2008-2012). The three global land cover maps describe all the terrestrial areas by 22 land cover 239 classes explicitly defined by a set of classifiers according to the United Nations Land Cover 240 Classification System, each classifier referring to vegetation life form, leaf type and leaf 241 longevity, flooding regime, non-vegetated cover types and artificiality (Di Gregorio, 2005).

242 The whole archive of full (300 m) and reduced resolution (1000 m) MERIS data acquired from 243 2003 to 2012 was first pre-processed and successfully fused as surface reflectance thanks to a set 244 of improved algorithms for radiometric calibration, geometric and atmospheric corrections, and 245 advanced cloud screening. A per pixel classification process, combining machine learning and 246 unsupervised algorithms, was then applied to the full time series to serve as a baseline to derive 247 land cover maps corresponding to each epoch. As temporal consistency was found as the most 248 important requirement for the climate modeling community, a multi-year integration strategy 249 was chosen for its better performance in reducing variability and improving stability (Bontemps 250 et al., 2012). Detected from the full-resolution Satellite Pour l'Observation de la Terre (SPOT) 251 vegetation time series (1998-2012), the land cover change corresponding to each epoch was 252 applied through back- and up-dating methods but only concerning the main macroscopic changes 253 observed for the forest classes (Li et al., 2016).

Inland open-water bodies and coastlines were mapped using wide-swath mode, image mode at medium-resolution (150 m) acquired by the Advanced Synthetic Aperture Radar sensor aboard ENVISAT satellite for a single period (2005-2010) (Santoro and Wegmüller, 2014) and then largely complemented with ancillary data.

The accuracy of the 2010 land cover product was estimated to 74.1% using the 2308 samples globally distributed and interpreted by regional experts. Further information on the accuracy of the ESA CCI land cover product in comparison to other existing global land cover data sets is provided by Tsendbazar et al. (2015).

In order to transform the ESA CCI land cover in Plant Functional Types (PFTs) distribution useable in ESMs, a CCI land cover user tool available from the visualization interface (http://maps.elie.ucl.ac.be/CCI/viewer/) can be used to apply a default or user-defined crosswalking table converting each land cover class into the corresponding proportions of PFT at the pixel level. This conversion also includes an aggregation of the different PFT distribution to coarser resolution grid cell in various projection systems.

In addition, the ESA CCI land cover products include information on the land surface seasonality at 1 km resolution which comprise climatological information of the vegetation greenness from Normalized Differenced Vegetation Index (NDVI) data as well as probabilities of snow and fire occurrences on a weekly basis at the pixel level. These were derived from SPOT vegetation daily observations from 1998 to 2012 as well from the corresponding MODIS time series. The interannual variability of these land surface seasonality variables was also computed from these 15year time series on a weekly basis that can be used for comparison with models. 275 **2.6** Aerosol

276 The ESA aerosol CCI team produces several long-term aerosol data sets (Popp et al., 2016) in 277 response to Global Climate Observing System (GCOS) requirements, including variables such as 278 aerosol optical depth (AOD) (from two Along-Track Scanning Radiometers (ATSR), the 279 MEdium Resolution Imaging Spectrometer (MERIS) and the POLarization and Directionality of 280 the Earth's Reflectances (POLDER) instrument), and stratospheric vertical extinction profiles 281 (using stellar occultation by the Global Ozone Monitoring by Occultation of Stars (GOMOS) 282 instrument). In response to the AEROCOM (http://aerocom.zmaw.de/) modeling community 283 needs, also information on aerosol composition such as fine-mode AOD (from radiometers) or 284 dust AOD (from the Infrared Atmospheric Sounding Interferometer IASI) and absorption AOD 285 (from ATSR and POLDER) are derived from the retrieved mixing ratio of various aerosol 286 components. All products include uncertainty estimates and are validated versus ground-based 287 reference data (AERONET, Holben et al., 1998) by independent experts. For the retrieval of 288 aerosol parameters from ATSR and IASI observations several algorithms are used, each of which 289 applies different physical principles and mathematical methods and thus different solutions to the 290 inversion problem. In the case of the ATSR radiometers, three algorithms (ADV from FMI, 291 ORAC from Oxford University and RAL and SU from Swansea University) do perform very 292 similarly, but with regional differences in both coverage and quantitative results, with none of 293 them performing better than the others everywhere (de Leeuw et al., 2015).

The ESA CCI aerosol product used in this paper is the 17-year climate data record including total AOD and fine mode AOD, both at 550 nm, produced by SU (version 4.21) using data from two similar sensors: the ATSR-2 on the European Remote Sensing Satellite 2 (ERS-2-ATSR-2), covering the time period 1995-2003, and the Advanced ATSR (AATSR) on ESA's 298 Environmental Satellite (ENVISAT-AATSR, from 2002 to April 2012). Level 3 (L3) monthly 299 mean data are used and only years with a full 12 months of data coverage are considered. 300 Incomplete years from either platform (1995, 1996 and 2003) are not taken into account, 301 restricting our analysis to the time period 1997-2011. The agreement of the data from the two 302 ATSR instruments during the overlapping period 2002-2003 was found to be very good making 303 it easy to combine the two data sets into a single time series. Here, we focus on total aerosol 304 optical depth (od550aer), fine mode AOD (od550lt1aer), absorption optical depth at 550 nm 305 (abs550aer) and AOD at 870 nm (od870aer). As an alternative observational data set, we use the 306 L3 collection 6 data from the Moderate Resolution Imaging Spectroradiometer (MODIS; Levy et 307 al., 2013) onboard Terra covering the time period 2003-2014.

308 **2.7 Ozone**

309 The ESA ozone CCI team produces a large number of L2 and L3 ozone data sets derived from 310 various satellite sensors operating in nadir, limb and solar/stellar occultation geometries (see e.g. 311 Miles et al., 2015; Lerot et al., 2014; Sofieva et al., 2013). In this work we use the total column 312 ozone (toz) data sets which consist of combined and harmonized L3 data covering the time 313 period between 1997 and 2010 (Coldewey-Egbers et al., 2015). Data from three 314 platforms/instruments, the Global Ozone Monitoring Experiment (GOME) onboard the European 315 Research Satellite 2 (ERS-2/GOME, 1996-2003), ENVISAT/SCIAMACHY (2003-2007), and 316 GOME-2 onboard the Metop satellites (METEOP/GOME-2, 2007-2011) are provided as a 317 merged gridded data set.

In additions to the total ozone data sets, we also include the ESA CCI limb gridded profile data, which consist of merged L3 monthly and zonally averaged data covering the time period 2007-2008 based on six different sensors, the MIPAS, SCIAMACHY, and GOMOS instruments onboard the ESA ENVISAT platform, the Optical Spectrograph and InfraRed Imaging System
(OSIRIS) and the Sub-Millimetre Radiometer (SMR) onboard of Odin, and the Atmospheric
Chemistry Experiment (ACE) instruments on Canadian SciSat platform.

324 The ozone CCI data sets used in this work have been extensively validated against ground-based 325 networks of Dobson and Brewer total ozone spectrophotometers (Koukouli et al., 2015), as well 326 as reference profile data sets from ozone sonde and lidar instruments (Hubert et al., 2016; 327 Keppens et al., 2015). These studies have demonstrated that CCI total column ozone data sets 328 closely match the accuracy and stability requirements defined by GCOS. Ozone profile data also 329 comply with GCOS requirements but only in a limited range of altitudes, covering the mid- to 330 upper stratosphere. In the upper-troposphere and lower stratosphere, the accuracy, precision and 331 stability of current data sets are still to be improved. Validation studies concentrating on L3 332 products have shown that the main source of uncertainty in gridded or merged data sets is related 333 to the limited sampling of satellite instruments. This source of uncertainty is especially 334 significant in polar spring conditions when the ozone field is characterized by a large variability 335 in space and time.

As an alternative reference data set for total ozone columns, we use data from the combined
NIWA data set (Bodeker et al., 2005) covering the time period 1980-2010.

338 **2.8** Greenhouse gases (GHG): XCO₂

The ESA CCI GHG product XCO_2 is retrieved from measurements of the two satellite instruments SCIAMACHY/ENVISAT (Bovensmann et al., 1999; Burrows et al., 1995) and TANSO-FTS/GOSAT (Kuze et al., 2009). XCO_2 is a dimensionless quantity (unit: ppm) defined as the vertical column of CO_2 divided by the vertical column of dry air (see Buchwitz et al. 343 (2005) for details). The XCO₂ distribution, the number of observations, the reported XCO₂
344 uncertainty and the XCO₂ standard deviation are available for 2003-2008 (land only) and 2009345 2014 (land and ocean).

346 XCO₂ is retrieved from radiance spectra in the near-infrared/short-wave infrared (NIR/SWIR) 347 spectral range using (mostly) optimal estimation (Rodgers, 2000) retrieval algorithms. Each 348 retrieval algorithm used to generate the corresponding Level 2 (L2) product has an underlying 349 radiative transfer model and a number of fit parameters (the so-called state vector elements), 350 which are iteratively adjusted until the simulated radiance spectrum gives an optimal fit to the 351 observed radiance spectrum (considering, e.g., instrument noise and a priori knowledge of 352 relevant atmospheric parameters). For details we refer to the Algorithm Theoretical Basis 353 Documents (ATBDs) available from the GHG-CCI website (http://www.esa-ghg-354 cci.org/sites/default/files/documents/public/documents/GHG-CCI DATA.html) for each 355 individual L2 data product. For the generation of the gridded L3 obs4MIPs product at monthly time resolution a spatial resolution of $5^{\circ}x5^{\circ}$ has been selected (instead of, e.g., $1^{\circ}x1^{\circ}$) to ensure 356 357 better noise suppression (note that the underlying individual satellite retrievals as contained in 358 the L2 products are sparse due to very strict quality filtering).

The gridded L3 obs4MIPs products have been generated from the individual sensor/algorithm L2 XCO₂ input data. In order to correct for the use of different CO₂ a priori assumptions in the independently retrieved products, all products have been brought to a common a priori using the Simple Empirical CO₂ Model (SECM) described by Reuter et al. (2012). After this, a gridded L3 product is generated from each L2 product by averaging all soundings onto a 5°x5° monthly grid. Only those grid cells are further considered having a standard error of less than 2 ppm. The grid cell uncertainty is computed from the reported L2 uncertainties and a term accounting for potential regional and temporal biases. To avoid potential discontinuities in the obs4MIPs time series, each L3 product has been offset corrected to have the same mean value of all overlapping grid boxes. The obs4MIPs XCO₂ value in a given grid cell is computed as the mean of the individual L3 values. Finally a filtering procedure has been applied to remove "unreliable" grid cells considering the overall noise error originating e.g. from instrumental noise (1.6 ppm) and total uncertainty (1.8 ppm) of each cell.

372 The obs4MIPs XCO₂ product has been validated by comparison with ground-based XCO₂ 373 retrievals from the Total Carbon Column Observation Network (TCCON, Wunch et al., 2011) 374 using version GGG2014 as a reference (Wunch et al., 2015). In short, the following has been 375 found: for XCO₂ the mean difference (satellite minus TCCON) is 0.3 ppm and the standard 376 deviation of the difference to TCCON is 1.2 ppm. The total uncertainty of the obs4MIPs product 377 is therefore about 1.5 ppm (1-sigma, per monthly 5°x5° grid cell, obtained via linear adding 378 instead of root-sum-square to be on the safe side). Details are given in Buchwitz and Reuter 379 (2016).

380 Due to the gridding / averaging process applied to generate obs4MIPs products detailed 381 time/location information is not available in the obs4MIPs data product and also averaging 382 kernels are not (yet) part of these products. Typically, however, the satellite XCO₂ averaging 383 kernel is close to unity. This is especially the case in the lower troposphere, where the CO₂ 384 variability is typically largest. Therefore applying the averaging kernels typically changes the 385 XCO₂ values by less than 1 ppm (Dils et al., 2014) and other error sources are likely more 386 relevant for using the obs4MIPs product such as the representativity error. A representativity 387 error originates from the fact that the GHG field from the obs4MIPs data set are derived by 388 averaging spatially and temporally sparse satellite observations, i.e., are not representative for the "true" monthly mean value of a given grid cell. Note that the validation results reported in the previous paragraph have also been obtained without considering the averaging kernels. The differences given above include to some extent the representativity error as well as other error sources such as the uncertainty of the TCCON reference observations, which is 0.4 ppm (1sigma). It is recommended to use the reported overall uncertainty range of 0.3 ± 1.2 ppm (1sigma) and/or the reported uncertainties for each grid cell as given in the obs4MIPs product file.

395 3 Models and simulations

In this study we use output from almost 50 global climate models (Table 1) that participated in
CMIP5 (Taylor et al. 2012). The model data were obtained from the World Climate Research
Programme's (WCRP) CMIP5 data archive made available through the Earth System Grid
Federation.

Table 1. CMIP5 coupled models used in this study (historical simulations extended beyond 2005 with RCP4.5 results). The models marked with asterisks (*) also provided model experiments with interactive ozone chemistry, models marked with daggers (†) also provided emission driven experiments with an interactive carbon cycle (historical emission driven simulations extended beyond 2005 with RCP8.5 results).

Model(s)	Host Institute	Resolution	References	
		(atmosphere)		
ACCESS1.0,	CSIRO (Commonwealth Scientific and	1.9°x1.3°, L38	Bi et al. (2013)	
ACCESS1.3	Industrial Research Organisation, Australia)			
	and BOM (Bureau of Meteorology,			
	Australia)			
BCC-CSM1.1,	Beijing Climate Center, China	2.8°x2.8°, L26;	Wu et al. (2010), Wu (2012)	
BCC-CSM1.1-M	Meteorological Administration, China	1.1°x1.1°, L26		
$BNU-ESM^{\dagger}$	College of Global Change and Earth System	2.8°x2.8°, L26	Ji et al. (2014)	
	Science (GCESS), BNU, Beijing, China			
CanCM4,	Canadian Center for Atmospheric Research,	2.8°x2.8°, L35	Arora et al. (2011)	
CanESM2 [†]	Canada			
CCSM4	National Center for Atmospheric Research	1.3°x0.9°, L26	Gent et al. (2011)	
	(NCAR), United States			
CESM1-BGC [†] ,	NSF/DOE NCAR (National Center for	1.3°x0.9°, L26;	Long et al. (2013)	
CESM1-CAM5,	Atmospheric Research) Boulder, CO, United	1.3°x0.9°, L26;	Hurrel et al. (2013)	
CESM1-CAM5-1-FV2,	States	1.3°x0.9°, L26;		
CESM1-FASTCHEM,		1.3°x0.9°, L26;		

CESM1-WACCM*		2.5°x1.9°, L66	Marsh et al. (2013)
CMCC-CM,	Centro Euro-Mediterraneo per i	0.8°x0.8°, L31;	http://www.cmcc.it/models;
CMCC-CMS	Cambiamenti Climatici (CMCC), Bologna, Italy	1.9°x1.9°, L95	Scoccimarro et al. (2011)
CNRM-CM5 [*]	Centre National de Recherches Météorologiques (CNRM), Météo-France and Centre Européen de Recherches et de Formation Avancée en Calcul Scientifique (CERFACS), France	1.4°x1.4°, L31	Voldoire et al. (2013)
CSIRO-Mk3.6.0	Australian Commonwealth Scientific and Industrial Research Organization (CSIRO) Marine and Atmospheric Research, Queensland Climate Change Centre of Excellence (QCCCE), Australia	1.9°x1.9°, L18	Rotstayn et al. (2010)
EC-EARTH	EC-Earth (European Earth System Model)	1.1°x1.1°, L62	Hazeleger et al. (2010)
FGOALS-g2,	Institute of Atmospheric Physics, Chinese	2.8°x3°, L26;	Li et al. (2010),
FGOALS-s2	Academy of Sciences (LASG)	2.8°x1.7°, L26	http://www.lasg.ac.cn/FGOA LS/CMIP5
$FIO-ESM^{\dagger}$	The First Institution of Oceanography, SOA, Qingdao, China	2.8°x2.8°, L26	Qiao et al. (2013)
GFDL-CM3 [*] ,	Geophysical Fluid Dynamics Laboratory	2.5°x2.5°, L48;	Donner et al. (2011);
$GFDL-ESM2G^{\dagger}$,	(NOAA GFDL), United States	2.5°x2°, L24;	http://nomads.gfdl.noaa.gov/
$GFDL$ - $ESM2M^{\dagger}$		2.5°x2°, L24;	
GFDL-CM2p1		2.5°x2°, L24	
GISS-E2-H [*] ,	Goddard Institute for Space Studies	2.5°x2°, L40	Schmidt et al. (2006)
GISS-E2-H-CC,	(NASA/GISS), United States	,	
GISS-E2-R [*] ,			
GISS-E2-R-CC			
HadCM3,	Met Office Hadley Centre, United Kingdom	3.8°x2.5°, L19;	Collins et al. (2001); Collins
HadGEM2-AO HadGEM2-CC.		1.9°x1.3°, L60;	et al. (2008); Collins et al. (2011)
HadGEM2-ES		1.8°x1.3°. L38	
INM-CM4	Institute for Numerical Mathematics (INM), Russia	2°x1.5°, L21	Volodin et al. (2010)
IPSL-CM5A-LR,	Institut Pierre Simon Laplace (IPSL), France	3.8°x1.9°, L39;	Dufresne et al. (2013),
IPSL-CM5A-MR.	1 ()/	2.5°x0.6°. L39:	Hourdin et al. (2013)
IPSL-CM5B-LR		3.8°x1.9°, L39	
MIROC-ESM [†] .	Japan Agency for Marine-Earth Science and	2.8°x2.8°. L80:	Watanabe et al. (2011):
MIROC-ESM-CHEM*	Technology (JAMSTEC) Atmosphere and	$2.8^{\circ}x2.8^{\circ}L80^{\circ}$	Sakamoto et al. (2012).
MIROC4h	Ocean Research Institute (AORI) University	$0.6^{\circ} \times 0.6^{\circ}$ L56	Watanabe et al. (2012) ,
MIROC5	of Tokyo and National Institute for	$1.4^{\circ}x1.4^{\circ}$ I 40	Watahabe et al. (2010)
Millious	Environmental Studies (NIES) Japan	1.1 XI.1 , E10	
MPI-FSM-LR [†]	Max Planck Institute for Meteorology	1 9°x1 9° I 47	Roeckner et al. (2006):
MPI-FSM-MR	Germany	1.9 AI.9 , EI7	10000kiler et ul. (2000),
MPI-FSM-P	Germany		Stevens et al. (2013)
MRI-CGCM3	Meteorological Research Institute (MRI)	1 1°x1 1° T 48	Yukimoto et al. (2013)
MRI-FSM1 [†]	Ianan	··· · ··· , 1.10	Turinoto et ul. (2011)
NorFSM1-M	Norwegian Climate Centre Norway	2.5°x1.9° L.26	Bentsen et al. (2013)
NorESM1-MF [†]	The wegan enhalt centre, the way	2.5 AL.7, 120	Bentsen et al. (2013)

For all variables except for column average CO_2 (XCO₂), we analyze the concentration driven CMIP5 historical simulations - twentieth-century simulations for 1850-2005 conducted with the best record of natural and anthropogenic climate forcing. In order to extend the model runs beyond the year 2005, we use results from simulations forced under the Representative Concentration Pathways 4.5 for the years 2006-2014. RCP4.5 is a scenario applied within

405 CMIP5 prescribing future greenhouse gas concentrations and resulting in a radiative forcing of 4.5 W m⁻² in the year 2100 relative to pre-industrial values (Clarke et al., 2007; Smith and 406 Wigley, 2006; Wise et al., 2009). The differences in the forcings between 2006 and 2014 for the 407 408 different emission scenarios (RCP2.6, RCP4.5, RCP8.5) are rather small and negligible 409 compared with the variability of the ensemble members of an individual model. We chose the 410 RCP for which the most data for the analyzed ECVs were available, which is RCP4.5.

411 For aerosol and ozone, the evaluation is only performed for the subset of CMIP5 models that has 412 interactive aerosols and chemistry, respectively.

Since CO_2 is prescribed in the concentration driven historical simulations, we analyze the 413 414 emission driven historical simulations (esmHistorical) for XCO₂. In this case, the simulations 415 were extended beyond 2005 with the corresponding RCP8.5 (esmrcp85) simulations because 416 emission driven simulations for RCP4.5 were not part of the CMIP5 experiment design.

417 If there are multiple ensemble members available for any given model, we only consider the ensemble member "r1i1p1" in our analysis. The only exceptions to this are the EC-EARTH 418 419 model, for which complete data sets were only available for "r6i1p1" and the GISS-E2-H and 420 GISS-E2-R models for which we used ensemble members with interactive ozone chemistry ("r1i1p2"; see section 5.7). 421

422

CMIP5 summary statistics 4

423 An assessment of the agreement of simulated climatological mean state and seasonal cycle for 424 key variables such as ECVs with observations is commonly seen as a reasonable starting point 425 for the evaluation of ESMs (e.g., Gleckler et al., 2008; Flato et al., 2013; Hagemann et al., 2013; 426 Eyring et al., 2016b). Following Gleckler et al. (2008) and similar to Fig. 9.7 of Flato et al.

427 (2013), we start the evaluation of the models by calculating the normalized relative space-time 428 RMSD of the climatological seasonal cycle from CMIP5 simulations compared with 429 observations for selected variables (section 4.1) and extended Taylor diagrams summarizing the 430 multi-year annual mean performance (section 4.2). For land use variables, no summary statistics 431 are calculated because the observations are rather static, i.e. do not provide a seasonal cycle.

432 All variables except for sea ice concentration are averaged over the whole globe. Sea ice 433 concentration is averaged over the latitude band 60°N to 90°N (Arctic, "NHpolar") and 60°S to 434 90°S (Antarctic, "SHpolar"). The model results are compared to a reference data set (marked 435 with asterisks in Table 2) and - where other data are available - to an alternative observationally 436 based data set. Table 3 gives an overview of the variables and the corresponding CMOR names 437 used while the observationally based data sets used for the evaluation are summarized in Table 2. 438 For the models, results are averaged over the years with observational data available given in 439 Table 2. Note that if alternative observationally based data are available, only years covered by 440 both, the reference and the alternative observations, are used.

Table 2. Observationally based data sets used for the model evaluation. The data sets marked with asterisks (*) are used as reference data sets in Figure 1 (lower right triangles), the other data sets are used as alternative data sets (upper left triangles in Figure 1, red stars in Figure 2). The variable names are defined in Table 3. The years specify the periods analyzed in Figure 1 and Figure 2.

Data set	Туре	Variable(s)	Resolution	Years (Figure 1, Figure 2)	Estimate of systematic errors	Reference(s)
AIRS_L3_RetStd-v5	satellite	hus	1°x1°	2003-2010	~25%	Tian et al. (2013), Susskind et al. (2006)
BDBP	ozonesond es	tro3	-	2006-2007		Hassler et al. (2008, 2009)
CERES-EBAF [*]	satellite	rlut, rsut, sw_cre, lw_cre	1°x1°	2001-2012	~5 W m ⁻²	Loeb et al. (2009, 2012)
CLARA-A2	satellite	clt	0.5°x0.5°	1982-2014		Karlsson et al. (2013),

						http://dx.doi.org/10.5 676/EUM_SAF_CM/ CLARA_AVHRR/V 002
ERA-Interim [*]	reanalysis	ta, tas, ua, va, zg;	0.75°x0.75°	1980-2005;		Dee et al. (2011)
		hus;		2003-2010;		
		clt		1982-2014		
ESA CCI Aerosol	satellite	od550aer; od870aer, od550lt1aer, abs550aer	1°x1°	2003-2011; 1997-2011		Popp et al. (2016)
ESA CCI Cloud [*]	satellite	clt	0.5°x0.5°	1982-2014		Stengel et al. (2016b)
ESA CCI Greenhouse Gases	satellite	xco2	5°x5°	2009-2014	~1.5 ppm	Buchwitz and Reuter (2016)
ESA CCI Ozone [*]	satellite	toz; tro3	1°x1°; 360°x10°	1997-2010; 2007-2008		Van Roozendael et al (2015)
ESA CCI Land Cover	satellite	lccs_class	300 m	2000, 2005, 2010		Defourny et al. (2015)
ESA CCI Sea Ice [*]	satellite	sic	25 km x 25 km	1992-2008		Sandven et al. (2015)
ESA CCI Sea Surface Temperature [*]	satellite- based/anal vsis	ts	0.05°x0.05°	1992-2010	~0.05 K (global median)	Merchant et al. (2014a,b)
ESA CCI Soil Moisture [*]	satellite	sm	0.25°x0.25°	1988-2005	$\sim 0.05 \text{ m}^3 \text{ m}^{-3}$	Liu et al. (2011, 2012), Wagner et al. (2012)
GPCP_L3_v2.2*	satellite + gauge	pr	2.5°x2.5°	1980-2005	0-2 mm day ⁻¹	Adler et al. (2003), Huffman and Bolvin (2013)
HadISST	satellite- based/anal vsis	ts	1°x1°	1992-2010		Rayner et al. (2003)
IGAG/SPARC	satellite + ozonesond es + model+ analysis	tro3	5°x5°	1960-2008		Cionni et al. (2011)
MODIS	satellite	clt; od550aer	1°x1°	2003-2014; 2003-2011		Platnick et al. (2003); Remer et al. (2005); doi: 10.5067/MODIS/MY
NCEP	reanalysis	ta, tas, ua,	2.5°x2.5°	1980-2005		D08_M3.006; Kalnay et al. (1996)
NIWA	satellite	toz	1.25°x1°	1997-2010		Bodeker et al. (2005)
NSIDC-NT, NSIDC- BT	satellite	sic	25 km x 25 km	1992-2008		Cho et al. (1996), Comiso and Nishio (2008), Comiso (1995)
PATMOS-x	satellite	clt	1°x1°	1982-2014		Heidinger et al. (2014)

Table 3. Variables used.

Variable	Name	Unit	Comment
abs550aer	Ambient aerosol absorption optical thickness at 550 nm	1	

clt	Total cloud fraction	%	For the whole atmospheric column, as seen from the surface or the top of the atmosphere; includes both large-scale and convective clouds
hus	Specific humidity	1	
lccs class	Land cover class	-	
lw_cre	Longwave cloud radiative effect	W m ⁻²	At the top of the atmosphere
mrsos	Soil moisture in upper portion of soil column	kg m ⁻²	Mass of water in all phases in a thin surface soil layer
od550aer	Ambient aerosol optical thickness at 550 nm	1	AOD from the ambient aerosols (i.e., includes aerosol water); does not include AOD from stratospheric aerosols if these are prescribed but includes other possible background aerosol types
od550lt1aer	Ambient fine aerosol optical thickness at 550 nm	1	od550 due to particles with wet diameter less than 1 µm ("ambient" means "wetted"); when models do not include explicit size information, it can be assumed that all anthropogenic aerosols and natural secondary aerosols have diameter less than 1 µm
od870aer	Ambient aerosol optical thickness at 870 nm	1	AOD from the ambient aerosols (i.e., includes aerosol water); does not include AOD from stratospheric aerosols if these are prescribed but includes other possible background aerosol types
pr	Precipitation	$kg m^{-2} s^{-1}$	At surface; includes both liquid and solid phases from all types of clouds (both large-scale and convective)
rlut	TOA outgoing longwave radiation	W m ⁻²	At the top of the atmosphere
rsut	TOA outgoing shortwave radiation	W m ⁻²	At the top of the atmosphere
sic	Sea ice area fraction	%	Fraction of grid cell covered by sea ice
sm	Volumetric soil moisture in upper portion of soil column	m ³ m ⁻³	Volume of water in all phases in a thin surface soil layer
sw_cre	Shortwave cloud radiative effect	W m ⁻²	At the top of the atmosphere
ta	Air temperature	К	
tas	Near-surface air temperature	K	
toz	Total ozone column	DU	Equivalent thickness at standard temperature and pressure (stp) of atmosphere ozone content
tro3	Ozone volume mixing ratio	ppbv	•
ts	Surface temperature	K	"skin" temperature (i.e., SST for open ocean)
ua	Eastward wind	$m s^{-1}$	
va	Northward wind	m s ⁻¹	
xco2	column average CO ₂ concentration	ppm	
zg	Geopotential height	m	Geopotential height

441 **4.1 Portrait diagram**

Figure 1 provides a synoptic overview of the relative quality of the CMIP5 models' representation of simulated climatological mean state and the seasonal cycle for ECVs compared with the multi-model median. The figure shows the relative space-time root-mean-square deviation (RMSD) from the climatological mean seasonal cycle assessing whether a specific model performs better or worse than the other models. The model data have been regridded to the grid of the reference data using a higher order patch recovery interpolation (Khoei and Gharehbaghi, 2007; Hung et al., 2004) and normalized with the centered median (i.e., subtracting the median and then dividing by the median). For the calculation of the RMSD, only grid cells with observational data available for at least 95% of the time period are taken into account.



Figure 1. Relative space-time root-mean-square deviation (RMSD) calculated from the climatological seasonal cycle of the CMIP5 simulations. The years averaged depend on the years with observational data available and are summarized in Table 2. A relative performance is displayed, with blue shading indicating better and red shading indicating worse performance than the median of all model results. A diagonal split of a grid square shows the relative error with respect to the reference data set (lower right triangle, data sets marked with asterisks in Table 2) and the alternative data set (upper left triangle). White

451

boxes are used when data are not available for a given model and variable. The variable names are defined in Table 3.

As such it can be seen as a starting point of model evaluation while the reasons for differences between model and observations need to be further investigated in additional analyses. The figure includes all variables that are shown in figure 9.7 of Flato et al. (2013) and adds variables with ESA CCI data now available. We would like to note that some differences compared to the portrait diagram of Flato et al. (2013) are introduced by using a different set of models, time range and observationally base reference data sets.

458 As found in previous studies, the performance varies across the models and variables, with some 459 models comparing better with observations for one variable and another model performing better 460 for a different variable. Typically, the multi-model mean outperforms any individual model, 461 which also holds for many of the newly added ECVs. Exceptions to this are, for example, global 462 average temperatures at 200 hPa (ta Glob-200), sea ice (sic NHpolar, sic SHpolar), aerosol 463 optical depth of fine particles at 550 nm (od550lt1aer Glob), and column average CO₂ 464 (xco2 Glob). In the following we discuss the results only for the variables that are compared to 465 the ESA CCI data sets and refer to Flato et al. (2013) for results on the other variables.

466 <u>SST:</u> typical biases in the geographical distribution of the simulated SST include a warm bias in 467 the subtropical stratocumulus regions as well as a cold bias in the equatorial Pacific. Individual 468 models performing worse than the multi-model mean (Figure 1) include, for instance, the 469 CSIRO, the FGOALS, and the MRI models. The reasons for this are rather different, for example 470 the CSIRO model shows a cold bias in the subtropical North Pacific whereas the FGOALS 471 model shows a warm bias in the subtropical Southeast Pacific. 472 <u>Sea ice:</u> for sea ice concentration (sic), the ESA CCI SI SSM/I and the National Snow and Ice 473 Data Center NSIDC-NT (Walsh et al., 2015) observations are used for comparison with the 474 CMIP5 models. Figure 1 shows that the choice of the reference data set does not impact the 475 results for the model performance in reproducing the observed sea ice concentration 476 significantly. This is expected as the two sea ice data sets are in rather good agreement.

477 Cloud: for total cloud cover (clt), the choice of the reference data set can make some difference 478 for the calculated performance of the individual models. A number of models such as, for 479 instance, the GFDL-CM3 and some of the HadGEM2 models have a larger RMSD when 480 compared against the ESA CCI data set than against the data from Pathfinder Atmospheres 481 Extended (PATMOS-x). The ESA CCI cloud data show slightly higher values (10-15%) for total 482 cloud cover in the subtropical stratocumulus regions off the west coasts of North and South 483 America as well as off the coast of Australia. In contrast, cloud amounts in the ESA CCI data are 484 smaller over the tropical Pacific with frequent deep convection (-10 to -20%). These are also 485 regions in which the models typically struggle to reproduce the observations. The average model 486 bias is therefore larger when the models are compared with the ESA CCI data rather than the 487 PATMOS-x data. An exact quantitative assessment, however, requires application of a satellite 488 simulator in the models to take into account satellite overpass times and lower cut-off thresholds 489 (Bodas-Salcedo et al., 2011), which is beyond the scope of this study. The comparison of total 490 cloud cover done here should therefore only be seen as a starting point for further evaluation of 491 the ESMs.

492 <u>Soil moisture:</u> the inter-model spread for soil moisture (sm) is large and most models tend to
493 systematically over- or underestimate soil moisture throughout the globe compared with the ESA
494 CCI data. It should be noted, however, that a quantitative comparison is difficult as the layer

thickness considered is not consistent among the models and the satellite observations (see
discussion in section 5.4). Qualitatively, many models such as the FGOALS, GFDL, HadGEM,
and MIROC models overestimate the soil moisture particularly in higher latitudes in Asia, as
well as Alaska and the northern part of Canada.

499 Aerosol: performance metrics for the four aerosol variables od550aer, od870aer, abs550aer, and 500 od550lt1aer are calculated with respect to the ESA CCI data set (see section 2.1) as reference 501 data set (lower triangles) and an alternative data set from MODIS. Shown are only CMIP5 502 models with interactive aerosols (ACCESS1-0, ACCESS1-3, BNU-ESM, CESM1-CAM5, 503 CSIRO-Mk3-6-0, GFDL-CM3, GFDL-ESM2G, GFDL-ESM2M, GISS-E2-H, GISS-E2-R, 504 HadGEM2-CC, HadGEM2-ES, IPSL-CM5B-LR, MIROC4h, MIROC5, MIROC-ESM, 505 MIROC-ESM-CHEM, MRI-CGCM3, NorESM1-M, NorESM1-ME), models using pre-scribed 506 aerosol climatologies have not been taken into account. Except for od550lt1aer, the multi-model 507 mean outperforms the individual models. Because of differences in the two satellite data sets for 508 AOD, which are largest over the continents (see section 5.1), the choice of the reference data set 509 can make a difference in the resulting model grading with most models performing slightly better 510 against MODIS than the ESA CCI data set. Additional analysis with the ESMValTool (not 511 shown) reveals that even though most models agree on the basic properties of the AOD 512 distribution (od550aer), the relative spread among the models for absorption AOD (abs550aer) 513 and AOD of fine particles (d < 1 μ m, od550lt1aer) is large. It should be noted that the 514 observational uncertainties for these quantities are also larger than for AOD at 550 nm. For 515 CMIP5, only the latter was evaluated whereas od550lt1aer, abs550aer, and od870aer are shown 516 for the first time here.

517 Ozone: the performance metric of total column ozone with respect to the ESA CCI (lower 518 triangle) and NIWA (upper triangle) data is shown only for models with interactive chemistry 519 (CESM1-WACCM, CNRM-CM5, GFDL-CM3, GISS-E2-H, GISS-E2-R, MIROC-ESM-520 CHEM). The performance of the individual CMIP5 models for global total column ozone is 521 quite similar for the two observational data sets. This is not surprising as both reference data sets 522 are based on the same satellite observations from GOME-2 and SCIAMACHY (Bodeker et al., 523 2005; Loyola et al., 2009). However, in the polar regions (toz SHpolar) there are significant 524 differences that likely occur because the ESA dataset has gaps in polar winter whereas these are 525 filled in the NIWA data set. Typical biases in CMIP5 models with interactive chemistry include, 526 for instance, an overestimation of total ozone in high northern latitudes (> $60^{\circ}N$) throughout the 527 year and an underestimation of ozone in Antarctica during summer (November to January) 528 (Eyring et al., 2013).

529 <u>CO₂</u>: only results from emission driven simulations are included in the performance metric 530 shown for XCO₂ in Figure 1. The BNU-ESM and the MPI-ESM-LR models outperform the 531 multi-model mean, which is biased high compared with the ESA CCI data as most models 532 systematically overestimate the column average CO₂ concentrations. This overestimation could 533 be possibly caused by slightly too weak CO₂ sinks in some models (Friedlingstein et al., 2014).

For most variables, the choice of reference data set does not make a big difference when using global averages for comparison with the CMIP5 models. This is, however, not necessarily the case when looking into more details such as individual regions or seasons. More on the comparison of the ESA CCI data with alternative observationally based data sets are given in the individual subsection of section 5.

539 4.2 Taylor diagrams

540 Another widely used way to summarize comparisons of results from a number of different 541 models with observations are Taylor diagrams (Taylor, 2001). The Taylor diagrams shown in 542 Figure 2 give the standard deviation and linear pattern correlation with observations of the total 543 spatial variability calculated from multi-year annual means, so in contrast to the space-time 544 RMSD evaluated in section 4.1, here only the geographical pattern is evaluated. For the calculation of the Taylor diagrams, all data have been regridded to a regular 1°x1° latitude-545 546 longitude grid using a patch recovery interpolation method. For each variable, a common 547 masking to exclude missing values has been applied to all data sets.

548 The standard deviations are normalized by the observed standard deviations, so the observed 549 climatology is represented in each panel by the filled black dots on the x-axis at x = 1. The 550 pattern correlation is given in this polar projection by the angular coordinate. The linear distance 551 between the observations and each model is proportional to the root-mean-square error (RMSE) 552 and can be estimated in multiples of the observed standard deviation with the gray circles 553 centered on the observational dots. The multi-model mean values have been calculated over all 554 models with data available (black star). Where available, an alternative reference data set (see 555 Table 2) is also shown in Figure 2 (red star).







Figure 2. Extended Taylor diagrams showing the multi-year annual average performance of the CMIP5 models in comparison with ESA CCI data for a) SST, b) total cloud cover, c) soil moisture, d) AOD at 550 nm, e) AOD at 870 nm, f) total column ozone, and g) column averaged CO₂ concentration. Panels a) to e) show CMIP5 historical simulations (extended with RCP4.5), panel f) historical simulations (extended with RCP4.5) with interactive ozone chemistry, and panel g) emission driven historical simulations (extended with RCP4.5). The multi-model mean values have been calculated over all models with data available (black stars). Where available alternative observationally based data sets are also shown (red stars, Table 2). The green circles show estimates of the observational uncertainties (RMSE, for details see section 4.2).

In this study, a new extended version of Taylor diagrams is presented that visualizes observational uncertainty: the green circles show estimates of the observational uncertainties (RMSE) that are part of the ESA CCI data sets. Here, the multi-year global average uncertainties given as one sigma of the total standard error normalized by the standard deviation of the observations are shown. The RMSE of a given model compared with the observations is therefore smaller than the 1-sigma uncertainty estimate of the observations if the model lies within the green circle.

564 <u>SST (Figure 2a):</u> the geographical annual mean patterns of the sea surface temperatures from the 565 models are highly correlated with the ESA CCI data with correlation coefficients ranging 566 between 0.94 and 0.98. However, SST in the subtropical stratocumulus regions as well in the 567 Southern Ocean is overestimated by many models. Another typical model bias found in many 568 simulations is an underestimation of the SST in the equatorial Pacific.

569 <u>Cloud (Figure 2b):</u> for total cloud cover, the models show a large spread in pattern correlation 570 between 0.25 and 0.88. Most models are, however, not outside of the 1-sigma uncertainty estimate showing that the differences between the models and the observations cannot be solelyexplained by model deficiencies.

573 <u>Soil moisture (Figure 2c):</u> can mostly be used for qualitative assessments of the models as the 574 observational uncertainties are larger than the RMSE of many of the individual models.

575 Aerosol (Figure 2d,e): the integrated aerosol properties AOD at 550 (Figure 2d) and 870 nm 576 (Figure 2e) also show a large inter-model spread. Because of the large observational 577 uncertainties, most models lie within the green circle of the 1-sigma measurement uncertainty 578 making further quantitative assessments difficult. This is also supported by the differences 579 between the ESA CCI data set and the MODIS data for AOD with MODIS being close to 1-580 sigma of the ESA CCI uncertainty estimate. The linear pattern correlation of most models with 581 the ESA CCI data, however, is smaller than that of the ESA CCI data and MODIS (0.8) showing 582 also differences in the geographical distribution of the simulated AOD (see also section 5.1 and 583 Figure 14).

<u>Ozone (Figure 2f)</u>: the correlation coefficients of the modeled total ozone columns with the ESA CCI data are quite high for most models (with interactive chemistry) with values above 0.94 and a ratio of the modeled and the observed spatial standard deviation close to 1. All models are, however, outside of the 1-sigma uncertainty estimate of the observations, which is also the case for the alternative observational data set (NIWA). Differences are found, for instance, in the northern high latitudes where the models tend to overestimate the total ozone columns (see also section 5.7 and Figure 18).

591 <u>CO₂ (Figure 2g):</u> For the column-averaged CO₂ concentrations, the correlation coefficients of the 592 results from the emission driven simulations with the ESA CCI data are typically quite low and

32

range between 0.4 and 0.6. This is partly caused by a systematical overestimation of XCO₂ concentrations by most CMIP5 models and partly by differences in the geographical patterns such as, for example, in northern Europe or Southeast Asia where the models show distinct local maxima that are not clearly visible in the ESA CCI data.

597 5 Further insights into the evaluation of CMIP5 models with ESA CCI data

In the following subsections, the evaluation of CMIP5 models using ESA CCI data and comparisons of ESA CCI data with alternative observational data sets are discussed individually for each of the CCI products (sea surface temperature, sea ice, cloud, soil moisture, land cover, aerosol, ozone, and greenhouse gases).

602 5.1 Sea surface temperature

The implemented diagnostics for sea surface temperature in the ESMValTool include the analysis of the temporal mean fields, their differences as well as a long-term trend analysis and calculation of scalar accuracy skill scores such as, for instance, area weighted RMSDs. All diagnostics can be applied to regional areas of interest defined by the user, e.g., ocean basins.

607 A major challenge when comparing the ESA CCI SST data to CMIP model results is that the 608 ocean grids used in the various CMIP models differ substantially. Thus, a common target grid 609 needs to be defined for the models and SST observations first. The user can specify the target 610 resolution and target projection in the ESMValTool configuration. For the examples given in Figure 3 and Figure 4, we use a T63 Gaussian grid as a common reference and project all SST 611 612 data to this grid using an energy conservative approach. In addition, the representativeness of the 613 SST variables largely varies among different models. While the CMIP sea surface temperature 614 variable (tos) corresponds to the temperature in a layer a few centimeters deep in some models, it represents the temperature of a layer of a couple of meters in other models. The ESA CCI SST product used in this study is designed to be representative of the sea surface at a depth of 20 cm. Except under conditions of very low wind stress and strong insolation, the stratification across the upper ~1 m of the ocean tends to be small because of near-surface mixing driven by wind and wave action. Nonetheless, the differing depth definitions need to be considered when interpreting SST differences between different models and between models and observations, particularly for the subset of the comparisons corresponding to situations of likely near surface stratification.

622 Figure 3 shows an example of a comparison of results from the CMIP5 model MPI-ESM-P with 623 the ESA CCI SST data set. On a global scale, the observed geographical patterns (top) with high 624 temperatures in the equatorial areas and low temperatures close to the poles are well reproduced 625 by the model and so is the global mean SST value of 287 K and its spatial variability. Both the 626 observations and the model show the typical two-armed warm areas in the Niño 3 and Niño 4 627 areas in the equatorial Pacific. They also show the typical shift of warm water in the eastern 628 northern Atlantic generated by the Gulf Stream, and the colder regions in the Arabian Sea. MPI-629 ESM-P shows a negative bias in the subtropics and tropics while a positive bias is found in the 630 cold climate zones in both hemispheres (Figure 3, bottom row). A switch in these differences 631 occurs in the temperate zones. The difference plot also shows discrepancies in specific areas 632 such as the underestimation of SST in the central northern Atlantic, from too-zonal behavior of 633 the North Atlantic Drift in the model, or the pattern of overestimation of temperature in ocean 634 upwelling zones on the east of ocean basins (along the Namibian coast, Baja California, etc.). 635 These discrepancies suggest differences in the representation of the wind driven upwelling and 636 western boundary currents.


Figure 3. Temporal means of SST in K for the ESA CCI data set (top right) and the CMIP5 model MPI-ESM-P (top left) as well as absolute (bottom left) and relative differences (bottom right).

638 Discrepancies between 7 exemplary CMIP5 models (GISS-E2-H-CC, GISS-E2-H, IPSL-CM5A-639 LR, MIROC-ESM-CHEM, MIROC-ESM, MPI-ESM-P, NorESM1-ME) for different ocean 640 basins are shown in Figure 4. Larger basins, like the northern or southern Pacific or Atlantic 641 Ocean, as well as the polar seas show good agreement in SST cycles among the different models 642 and with the ESA CCI data. Differences are larger for smaller basins like the Baltic or 643 Mediterranean Seas or the Niño regions. These larger discrepancies occur due to the size of these 644 smaller regions and their higher sensitivity to small scale fluctuations. Spatial averages of larger 645 basins attenuate such fluctuations. The ESA CCI SST data are sufficient to show the model 646 limitations on such scales, and discriminate, for example, the better seasonal cycle amplitude for 647 the Baltic Sea in MIROC-ESM-CHEM compared to MPI-ESM-P.



Figure 4. Time series of mean SST for different ocean basins from 7 CMIP5 models (see legend) compared with the ESA CCI SST data.

ESA CCI SSTs are relatively unusual in being physics-based (not tuned to drifting buoys) and explicitly aiming to represent the 20-cm depth SST, which should correspond well to modellayer-average SSTs in most circumstances. The new data set therefore provides an independent, accurate (0.1 K), high-stability climate data record.

653 **5.2 Sea ice**

The observed rapid decline in Arctic sea ice thickness and extent over the last few decades is one of the most striking indicators for climate change (Stroeve et al., 2012; Lindsay and Schweiger, 2015). The melting of sea ice contributes to the rise of global temperatures through the icealbedo feedback (Curry et al., 1995). The decline in sea ice extent is a positive feedback where the initial shrinkage in the area of sea ice reduces the albedo and thus reinforces the initial alteration in sea ice area. High-quality observations of sea ice are thus crucial to monitor climate change and to evaluate climate models.

661 Here, we use data from the National Snow and Ice Data Center (Walsh et al., 2015) as an 662 additional reference data set for the model evaluation and for comparison with the ESA CCI sea 663 ice data. The NSIDC provides two different data sets, each covering the time period from 1979 664 to present. The main difference between the two data sets is the algorithm used in processing the 665 satellite data: the NASA Team (NSIDC-NT; Cavalieri et al., 1996) and the Bootstrap (NSIDC-666 BC; Comiso, 2000) algorithm. While the NSIDC-BT algorithm corrects for melt ponds that are 667 treated as open water by synthetically increasing the summer sea ice concentration (sic), such a 668 correction is not included in NSIDC-NT.

669 The sea ice diagnostics implemented into the ESMValTool include time series of the modeled 670 and observed evolution of sea ice extent (Figure 5) or area as well as polar-stereographic contour 671 plots of sic and sic biases (Figure 6). The sea ice extent has been calculated by adding up the 672 surface area of all grid cells with a sea ice concentration equal or larger than 15%. Satellites in 673 polar orbits do not pass directly over the poles. As a consequence, there is a small area centered 674 around the poles that cannot be observed by these instruments. For the comparison with the 675 model data shown in Figure 5, these pole holes have been filled assuming 100% sea ice cover in 676 this region.

677 The time series of September Arctic sea ice extent in Figure 5 shows that the spread between the 678 four observational data sets (thick black lines) from ESA CCI and NSIDC is much smaller than 679 the spread among the CMIP5 models (colored lines), which amounts to about 9 million km^2 680 between CSIRO-Mk3-6-0 (largest positive bias) and GISS-E2-H (largest negative bias). 681 However, the CMIP5 multi-model mean (thick red line) lies most of the time within the 682 observational spread although the RCP4.5 simulation mean does not show the decrease in sea ice 683 extent that has been observed between 2005 and 2013. The sea ice extent from the ESA CCI data 684 sets (thick black lines) is in very good agreement with the NSIDC data sets. ESA CCI SSM/I data show a small positive bias compared with NSIDC-NT of up to 1 million km² between 1997 685 686 and 2005. ESA CCI AMSR-E data are in very good agreement with both NSIDC data sets. The negative trend over the observed time period from 1990 to 2010 is about 1 million km² per 687 688 decade in all four observational data sets. The magnitude of this trend is, however, 689 underestimated by the CMIP5 multi-model mean.



Figure 5. Evolution (1960-2020) of September Arctic sea ice extent in million km² from the CMIP5 models (colored lines) and from observations (thick black lines). The pole holes of the satellite data sets have been filled assuming a sea ice concentration of 100%. All available ensemble members from a given model are shown and drawn in the same color as indicated in the legend. The CMIP5 multi-model mean is shown in bold red and the gray shading shows the standard deviation of the CMIP5 ensemble. The observations are from ESA CCI SI and NSIDC. Figure modified from Bräu (2013).

690

Figure 6 shows polar-stereographic contour maps of Arctic September (upper row) and AntarcticMarch (lower row) sic, which roughly corresponds to the average annual minimum sea ice

693 extent. As in Figure 5, there is good agreement between the ESA CCI SI SSM/I (left column) 694 and NSIDC-NT (middle column) also in the geographical distribution. The sic from the two data 695 sets differs by less than 0.2 in all grid cells for both Arctic and Antarctic sea ice distributions (not 696 shown). In the Arctic, the CMIP5 multi-model mean slightly underestimates the observed sic in 697 the marginal ice zone of the Central Arctic Ocean and in the East-Siberian and Beaufort Seas by 698 about 0.2 (right column). There is also a small overestimation east of Svalbard. In the Antarctic, 699 the sea ice concentration is underestimated by the CMIP5 multi-model mean in the Weddell Sea 700 as well as in a belt along the coast of the Amundsen, Ross and Somov Seas by up to 0.6.



Figure 6. Polar-stereographic map of Arctic September (upper row) and Antarctic March (lower row) sea ice concentration from ESA CCI SI SSM/I (left column) and NSIDC-NT (middle column) observations

701

averaged over the years 1992-2008. The pole holes of the satellite data sets have been filled assuming a sea ice concentration of 100%. The right column depicts the differences between the CMIP5 multi-model mean and the ESA CCI SI SSM/I observations averaged over the years 1992-2005.

702 In general, the ESA CCI data show good agreement to the data sets from the NSIDC. For robust 703 assessments of retrospective climate simulations, however, a longer time period is needed and 704 would ideally go from the early 1980s to present. Since the sea ice observational data are no 705 exception in having errors that are inherent to all observations, the daily uncertainty estimates 706 provided by the ESA CCI sea ice team are very useful for a more quantitative model evaluation. 707 These error estimates are based on the extensive algorithm comparison study (Ivanova et al., 708 2015) and have been underpinned by subsequent validation studies (Kern et al., 2016). The error 709 estimates will be useful for further regional and seasonal assessments of sea ice concentrations.

710 **5.3 Cloud**

711 Clouds strongly affect the Earth's radiative balance and temperature but are challenging to model 712 and observe, leading to large uncertainties in understanding climate variability and change. 713 Model evaluation using long-term, consistent observational data records can help to improve 714 both, the understanding of the present-day climate and the confidence in climate model 715 projections. Modeled clouds and satellite observations are difficult to compare because observed 716 clouds are affected by the satellite instrument's sensitivity, the temporal and spatial sampling and 717 the vertical overlap of the cloud layers, while the clouds in climate models are assumed to be 718 plane-parallel and are of coarse horizontal and vertical resolution. Ideally, a satellite simulator 719 (e.g., Bodas-Salcedo et al., 2011) is used during the model simulation to mimic the satellite 720 viewing geometry, temporal sampling and specific instrument characteristics such as lower cut-721 off values. Many CMIP5 historical and future scenario simulations, however, have been run without such a satellite simulator. Total cloud cover is the model cloud parameter that most readily can be compared directly to the satellite derived cloud fraction without a simulator, even though models can have substantial cloud cover but very little cloud condensate making those clouds too optically thin to be detected by the satellite instrument.

726 Here we use the ESMValTool diagnostics mean, bias and interannual variability to compare 727 Cloud cci AVHRR-PM total cloud cover with other satellite-based cloud data sets and to 728 evaluate CMIP5 models. Figure 7 shows the ESA CCI total cloud cover (clt) in boreal winter 729 (December, January, February) and summer (June, July, August) and the associated total 730 uncertainties derived from comparisons with CALIOP as described in section 2.3. The inherent 731 AVHRR difficulties in detecting clouds during polar night and over high elevation, snow 732 covered areas (North Canada, North East Asia and Himalayas) result in uncertainties of more 733 than 20% in these regions. Comparing the ESA CCI zonal mean cloudiness to other AVHRR 734 cloud data sets such as PATMOS-x (Heidinger et al., 2014) and CLARA-A2 (Karlsson et al., 735 2013) and the MODIS cloud data set (Platnick et al., 2003) also show the largest observational 736 spread (40-50%) in high latitudes in the winter hemisphere. The ESA CCI uncertainties are also 737 high with values of up to 20% in the subtropical high pressure dry areas. In these regions, the 738 ESA CCI data set has 5-10% less cloud coverage than PATMOS-x and CLARA-A2 (not shown).

The performance metrics results in Figure 1 show that the cloudiness of most CMIP5 models compare well with the ESA CCI data and the alternative reference data set PATMOS-x on a global scale, but there are regional differences as seen in Figure 7. The CMIP5 multi-model mean bias compared to the ESA CCI data shows an underestimation of cloud amount especially in the subtropical stratocumulus regions off the west coasts of North and South America as well as off the coast of Australia as known from many previous studies (e.g., Nam et al., 2012). In 745 contrast, the CMIP5 multi-model mean and most individual models overestimate cloud amounts 746 by 20% over the subtropical high pressure regions with minimum cloud amounts. These biases 747 are smaller (10-15%) if the models are compared to PATMOS-x and CLARA-A2 instead 748 because cloud amounts from these two alternative observational data sets are larger than from the 749 ESA CCI data set in these regions. The CMIP5 models with a normalized RMSD above 0.2 in 750 Figure 1 (CCSM4, CESM1-BGC, HadCM3, MIROC-ESM and MIROC-ESM-CHEM) 751 underestimate cloud amount on a global scale (not shown). The largest inter-model spread (60%) 752 occurs at high latitudes in polar winter, where also the observational data sets have their largest 753 uncertainties as seen in the zonal mean plots in Figure 7. In these cold conditions the amount of 754 cloud condensate is small and the modeled clouds are often thinner than the satellites' detection 755 limit. Here, using a simulator removes part of these model clouds.



Figure 7. Maps of the multi-year seasonal mean of total cloud cover and 1-sigma uncertainty from ESA CCI cloud for a) December-January-February (DJF) and b) June-July-August (JJA) averaged over the years 1982-2014. The figure also shows the differences between the ESA CCI data and the CMIP5 multi-model mean as well as zonal means. The zonal mean panels show averages from ESA CCI (red), PATMOS-x (blue), CLARA-A2 (cyan), MODIS (green), ERA-Interim (orange), and the CMIP5 multi-model mean (black). The individual CMIP5 models are shown as thin gray lines and the observational uncertainties of the ESA CCI data (±1-sigma) are shaded in light red. The MODIS data are only available for the years 2003-2014.

757 Figure 8 shows the interannual variability of total cloud cover for the satellite data sets, the 758 CMIP5 multi-model mean and ERA-Interim. The interannual variability is estimated as the 759 relative temporal standard deviation of the deseasonalized monthly mean time series (Lauer and 760 Hamilton, 2013). All the AVHRR data sets (ESA CCI, CLARA-A2, PAMOS-x) have their 761 largest variability (30-40%) for the dry tropical high pressure regions over the oceans, over north 762 Africa, south Africa and Australia, reflecting the annual shift of the ITCZ and the El 763 Niño/Southern Oscillation (ENSO). MODIS tropical Pacific Ocean variability is smaller than in 764 the AVHRR data sets, since MODIS data are available only for the time period 2003-2014 and 765 thus do not include the strong El Niño events in the 1980s and 1990s, which illustrates the 766 importance of using long-term observational records when evaluating ENSO. The ESA CCI data 767 have larger variability over the tropical Pacific Ocean than the other AVHRR satellite data sets. 768 Time series (not shown) reveal that the ESA CCI clt is of similar magnitude as PATMOS-x and 769 CLARA-A2 for El Niño years when the cloud cover is maximum, while the ESA CCI data have 770 less cloud amount (5-15%) for La Niña years when the cloud cover reaches minima. This results 771 in a larger interannual variability of the ESA CCI data. PATMOS-x data show less variability and higher cloud amounts over the Antarctic than the ESA CCI and CLARA-A2 data. The CMIP5 multi-model mean shows less variability than the observations, especially over the subtropical high pressure regions, where most of the individual CMIP5 models overestimate the total cloud cover. In contrast, the models that underestimate clt in the dry regions (CCSM4, CESM1-BGC, HadCM3, MIROC-ESM, MIROC-ESM-CHEM) show a larger interannual variability.



Figure 8. Interannual variability in total cloud cover estimated from the relative temporal standard deviation of the deseasonalized monthly mean time series from 1982 to 2014. Shown are (from top left to bottom right) satellite data (ESA CCI cloud, CLARA-A2, PATMOS-x, MODIS) in comparison with ERA-Interim reanalysis data (lower row, center) and the CMIP5 multi-model mean (lower row, right). The MODIS data are only available for the years 2003-2014.

The Cloud_cci AVHRR-PM total cloud cover data compare well with other existing long-term AVHRR cloud data sets. The ESA CCI pixel-based uncertainties show the user which areas should be interpreted carefully, e.g. polar and high elevation snow covered regions where the passive satellites have problems detecting clouds. The ESA CCI cloud cover data show lower 783 minima than the other AVHRR data sets for the tropical Pacific, which should be investigated 784 further. The other ESA CCI cloud data sets with shorter time records (MODIS, ATSR-2, 785 AATSR and MERIS) can be used for process studies and for narrowing the observational 786 uncertainties. A Cloud cci satellite simulator has been developed, which can be used in future 787 CMIP simulations and include other cloud variables such as cloud top pressure, optical 788 thickness, effective radius, albedo and liquid/ice water path in the model evaluation. Cloud cover 789 from the CMIP5 models shows the known typical error patterns compared with the ESA CCI 790 data and the other satellite data sets, underestimating clouds in the stratocumulus regions and 791 overestimating clouds in the subtropical dry regions. More detailed analysis of the individual 792 models and the interaction with radiation are needed to understand these biases.

793 **5.4 Soil moisture**

794 The current soil moisture diagnostics implemented in the ESMValTool comprise metrics for the 795 evaluation of soil moisture from regional to global scale and are largely based on Loew et al. 796 (2013) using version 2.2 of the ESA CCI soil moisture data set. These include the comparison of 797 temporal mean fields of soil moisture, as well as the analysis of the co-variability of soil 798 moisture anomalies with precipitation anomalies and the similarity of the spatial patterns of the 799 percentile distributions of the model and observations. The latter is a measure for the similarity 800 of the spatio-temporal dynamics of the soil moisture field (see Loew et al., 2013 for details). 801 Another diagnostic analyzes the long-term trend in soil moisture for both the ESA CCI data set 802 and CMIP models. The non-parametric Mann-Kendall regression is used to assess the statistical 803 significance of long-term soil moisture trends, similar to Dorigo et al. (2012) for the time period 804 1988-2008. This time period was chosen because the ESA CCI soil moisture data have a poorer temporal sampling prior to this period (Loew et al., 2013). All diagnostics can be applied at theglobal scale as well as for user-defined regions.

807 A general challenge when comparing satellite soil moisture with model results is that the 808 observations represent a rather different quantity than the one simulated by the models. CMIP 809 models provide the soil moisture as storage terms for soil layers at specific depths. As the 810 different CMIP models are based on different soil model implementations, these are not 811 necessarily directly comparable as they might differ in their depth and therefore in their temporal 812 dynamics. Currently, the official CMIP5 output comprises two soil moisture variables, which are 813 supposed to represent a 10-cm surface layer (mrsos) or the entire soil column (mrso). Here, we 814 use only data from models that provided the surface layer soil moisture for comparison. The 815 surface layer soil moisture is converted into the volumetric soil moisture content by dividing 816 mrsos by the thickness of the represented layer and by the density of water, which is assumed to be 998.2 kg m⁻³ (20°C). The variable for volumetric soil moisture content compared with the 817 818 ESA CCI data is called sm (see Table 3).

Satellite soil moisture data typically represent the volumetric soil moisture content (m³ m⁻³) of a shallow surface layer, which is also the case for the ESA CCI data set. The soil moisture diagnostics implemented in the ESMValTool compare the volumetric soil moisture content calculated from the model output with observations. All data are aggregated to similar temporal and spatial scales before further analysis.

Figure 9 shows an example of the ESA CCI volumetric soil moisture data compared with soil moisture from the CNRM-CM5 model. The model shows comparable soil moisture patterns in large parts of the globe. A wet bias is observed in the northern latitudes, which might be related 827 to an overestimation of soil moisture due to missing processes in the model (e.g., freeze-thaw 828 dynamics). The model bias can also be related to a dry bias in the ESA CCI observations in these 829 regions as no soil moisture is observed during wintertime and under frozen soil conditions. 830 Relative differences are largest in the desert regions (Sahara, Arabian Peninsula), which is, 831 however, of minor importance due to the overall small absolute soil moisture content in these 832 regions. The wet region along the southern border of the Himalayas is clearly visible in the 833 model but not in the ESA CCI soil moisture. This is most likely due to the complexity in 834 mountainous terrain with large terrain slopes (for both the models and in the satellite soil 835 moisture retrieval algorithms).



Figure 9. Temporal mean fields of volumetric soil moisture from the CNRM-CM5 model (top left), the ESA CCI soil moisture data set (top right) as well as their absolute (bottom left) and relative differences (bottom right).

836

The long-term trends in soil moisture during the time period 1988-2008 are compared in Figure 10. The figure illustrates only statistically significant trends (p < 0.05). The ESA CCI soil moisture shows decreasing soil moisture in large parts of the globe. Strongest decline of soil moisture is observed in southern Russia, while positive trends are observed in the tropical parts of Africa. Trends in the CNRM-CM5 model are rather different to those obtained from the CCI data set. A significant decline in soil moisture is observed over Europe, while a significant increase of soil moisture is simulated throughout large parts of the northern hemisphere.



Figure 10. Temporal trend in soil moisture over the period 1988-2008 as derived from the CNRM-CM5 model (left) and the ESA CCI soil moisture data sets (right). Masked areas represent grid cells where the Mann-Kendall correlation coefficient was not statistically significant at the 95% confidence level.

The percentiles of the observed and simulated soil moisture fields are rather similar, which illustrates that both data sets show similar spatial patterns of the soil moisture dynamics. As an example, Figure 11 shows the percentile maps for the 5%, 50% and 95% percentiles for the observed and simulated (CNRM-CM5) soil moisture fields. For each of the percentiles, the spatial autocorrelation coefficient results in very high correlation values ($\rho > 0.9$), which indicates a strong similarity of the spatial patterns. Percentile: 0.05 (r=1.0)



851

Figure 11. Percentile maps for ESA CCI soil moisture (left column) and soil moisture from CNRM-CM5 (right column). The (from top to bottom) 5%, 50% and 95% percentiles are shown and the spatial correlation coefficient between the model and the observations is provided in the title of each plot.

There is increasing evidence on the quality and consistency of the trends in the ESA CCI soil moisture data set. For example, in a recent special issue in the International Journal of Applied Earth Observation and Geoinformation (JAG) (vol. 48, June 16) several trend papers are presented and reveal reliable trends over many parts of the globe where they were compared with other water related observations including runoff, precipitation, and reanalysis data (see e.g., Wang et al., 2016; Su et al., 2016; Du et al., 2016; Qiu et al., 2016, all in the special issue in JAG). These results give more confidence in the ESA CCI soil moisture trends, especially over the sparsely to moderately vegetated regions. This is also highly relevant to assessing soil moisture variability and change in the context of a changing climate, which has been a great challenge so far.

862 **5.5 Land cover**

863 Benchmarking climate models with land cover information is not straight forward due to the 864 different concepts of representation of terrestrial vegetation in global Dynamic Vegetation 865 Models (DGVM), which are typically based on the concept of PFTs that are supposed to 866 represent groups of land cover with similar functional behavior. Thus, an important first step is 867 to map the ESA CCI land cover classes to PFTs like the ones used in CMIP models (Figure 12) 868 (Poulter et al., 2015). As the PFTs in CMIP models differ, the current ESMValTool diagnostics 869 analyzes only broad surface types (bare soil, grass, shrubs, forests), which is similar to the 870 approach chosen by Brovkin et al. (2013). Land cover is either prescribed in the CMIP models or 871 simulated using a DGVM. In particular for the latter case, an independent assessment of the 872 accuracy of the simulated spatial distributions of major land cover types is desirable in order to 873 evaluate the DGVM accuracy for present climate conditions. The diagnostic currently 874 implemented into the ESMValTool considers the land cover to be static for present climate 875 conditions in the CMIP models. The PFT distribution is then compared against satellite 876 observations from a similar time period.



Figure 12. Area fraction (%) of forest and shrub cover in the MPI-ESM-MR model (top left) and the ESA CCI land cover data set (top right) and absolute (bottom left) and relative differences (bottom right). The ESA CCI 2005 epoch was used for the analysis.

877

Figure 12 and Figure 13 show differences in the area cover fraction for forest type land covers as well as grassland and cropland areas between the ESA CCI land cover product and the MPI-ESM-MR model, which is based on the DGVM JSBACH for the terrestrial component (Brovkin et al., 2009; Brovkin et al., 2013). The tree cover in MPI-ESM-MR is underestimated compared to the ESA CCI data set in the Amazon and along the west coast of North America, while grass and cropland is overestimated in many parts of the globe. Similar analysis results are obtained when using the ESA CCI epoch for the year 2000 instead of 2005.



Figure 13. Area fraction (%) of grass and cropland cover in the MPI-ESM-MR model (top left) and the ESA CCI land cover data set (top right) and absolute (bottom left) and relative differences (bottom right). The ESA CCI 2005 epoch was used for the analysis.

The ESA CCI land cover data set provides the first consistent series of high-resolution (300 m) global land cover products derived by combining a whole suite of different sensors including information on PFTs. This has become important in particular for evaluation of ESMs that start to include more complex land cover dynamics in projections of future climate.

890 **5.6 Aerosol**

885

The geographical distribution of the multi-year averages of od550aer, o550lt1aero, and abs550aer, as well as the differences between the ESA CCI data and some exemplary CMIP5 models (CSIRO-Mk3-6-0, GFDL-CM3, GISS-E2-H, IPSL-CM5B-LR, MIROC-ESM-CHEM) are shown in Figure 14. Here, we consider only the CMIP5 models with interactive aerosols and

895 exclude multiple versions of the same model. In general, the models' performance is better over 896 the oceans than over the continents although the SU algorithm used to process the CCI data may 897 underestimate AOD over the oceans. Large model biases are found over the Sahara where some 898 models (especially GFDL-CM3 and IPSL-CM5B-LR) underestimate the aerosol optical depth 899 (left column). This could be caused by an incorrect representation of dust which is consistent 900 with a much better performance of these models for the fine mode optical depth (middle column) 901 in the same region. In addition, the underestimation of AOD over the Sahara might also be partly 902 amplified by an overestimation of AOD in the ESA CCI aerosol product (SU), which is a known 903 problem in this region. In contrast, a substantial positive bias is found over Europe and East Asia 904 (in particular CSIRO-Mk3-6-0 and GISS-E2-H) with similar biases both in the total and in the 905 fine mode optical depth. Significant deviations from the observations are also visible in the 906 modeled absorption optical depth (right column), especially in tropical regions. The contribution 907 of absorption to the aerosol optical depth is, however, quite small. We also note that the satellite 908 uncertainty for abs550aer is larger than for od550aer and od550ltaer.



Figure 14. Climatological mean AOD (left column), fine mode optical depth (middle) and absorption optical depth (right column) at 550 nm averaged over the period 1997-2011. The first row shows the the observations (ESA CCI ATSR SU v4.21), the other rows the differences between selected CMIP5 models with interactive aerosols and the ESA CCI data. Differences that are not statistically significant at the 95% confidence level are masked out in gray.

910 As can also be seen in Figure 1, the two satellite data sets used as observational references result 911 in different model performance grades, mainly because of measurement uncertainties inherent to 912 the data sets. To further explore the reason for these differences, the two satellite data sets are 913 compared with ground-based measurements from the AErosol RObotic NETwork (AERONET; 914 Holben et al., 1998) (Figure 15). AERONET data are widely accepted as a reliable reference for 915 aerosol optical depth and are often used for validating satellite products. AERONET data, 916 however, do not provide global coverage with very few measurements particularly over the 917 ocean. The few AERONET sites that are measuring AOD over the ocean are typically near 918 shallow-water areas such as on islands and the coastlines of continents, and thus not 919 representative of open ocean conditions. The Marine Aerosol Network MAN has therefore been established to provide AOD measured with hand-held sun photometers, predominantly on 920 921 research ships, starting from 2004 (Smirnov et al., 2009). However, in spite of the many cruises 922 included, the data are still sparse making global satellite data sets very valuable for evaluation of 923 ESMs. For consistency, we only consider years that are covered by both, the MODIS and the 924 ESA CCI data sets (2003-2011). Similarly to the models, the largest differences between the two 925 satellite data sets are found over the continents (top row of Figure 15). This is not surprising 926 given that satellite retrievals over the dark ocean surfaces are less sensitive to the assumptions in 927 the retrieval algorithms. The ESA CCI product shows a considerably higher optical depth than MODIS over the Sahara and seems to be in slightly better agreement with AERONET in this region (however only a few stations are available around the Sahara). Another striking difference between the two data sets is found over Southeast Asia where od550aer from MODIS is higher than the values from the ESA CCI resulting in a slightly better performance when compared to AERONET. The overall performance of the two data sets is quite similar but the MODIS data show a higher correlation ($R^2 = 0.85$) with AERONET than the ESA CCI data ($R^2 = 0.76$) as can be seen in the scatter plots in the bottom row of Figure 15.



Figure 15. Comparison of AOD at 550 nm from the ESA CCI ATSR SU v4.21 and the MODIS Terra C6 satellite products against the AERONET ground-based measurements for the period 2003-2011. The top

935

row shows the AERONET values as open circles plotted on top of the satellite data averaged over the same time period. The bottom row shows scatter plots of spatially and temporally collocated measurements on a monthly-mean basis.

With the ESA CCI aerosol and the MODIS data, two independent, long-term satellite data sets are available for model evaluation. This is particularly helpful when there is doubt about the reliability of the comparison with model results by adding the possibility to provide an independent check whether the satellite data are correct. Furthermore, in some areas, the ESA CCI aerosol products provide better correlation with AERONET than MODIS and the addition of ATSR to MODIS data can improve the overall results when used for data assimilation as there are more data available to constrain the model.

943 **5.7 Ozone**

For the first time in CMIP, a subset of the models included interactive chemistry in CMIP5. Also in contrast to previous CMIP phases, the models that prescribed ozone in CMIP5 included a time-varying stratospheric ozone climatology (Cionni et al., 2011) rather than a constant forcing. Detailed information on the treatment of ozone in CMIP5 models as well as an evaluation of their performance compared to observations is given in Eyring et al. (2013). Here we repeat some of this analysis by adding the newly available ESA CCI ozone data.

Eyring et al. (2013) divided the CMIP5 models into three classes: (a) CMIP5 models with interactive chemistry, (b) CMIP5 models with semi-interactive chemistry including those models that prescribed ozone data based on results from the underlying CMIP5 chemistry-climate model, and (c) CMIP5 models that prescribed ozone IGAC/SPARC ozone database (Cionni et al., 2011). Here, we focus on the models with interactive ozone chemistry only. The performance 955 of the individual CMIP5 models with interactive chemistry for total ozone columns is similar 956 with respect to both observational data sets (ESA CCI and NIWA) as can be seen in the time 957 series from 1960 through 2010 shown in Figure 16. Differences in both data sets are therefore 958 mostly a result of different statistical methods used to combine the different satellite data sets.

Most CMIP5 models with interactive chemistry overestimate the annual global mean total column ozone compared with the ESA CCI data (Figure 16a) but capture the trend of ozone depletion starting in the 1980s quite well. The October mean total column ozone in the Antarctic (90°S-60°S) is well captured by the CMIP5 models in terms of both, magnitude and trend (Figure 16b).



Figure 16. Time series of area-weighted total column ozone from 1960 to 2010 for a) global annual mean (90°S-90°N) and b) Antarctic October mean (60°S-90°S). The figure shows the multi-model mean (black line) and standard deviation (gray shading) as well as individual CMIP5 models with interactive chemistry (colored lines) compared with ESA CCI (filled circles) and NIWA (open triangles) data. The

IGAG/SPARC ozone database (Cionni et al., 2011) is also shown as a reference (orange line). All data sets have been interpolated to the same grid as the ESA CCI observations. During the periods covered by observations, only grid cells in the time series with valid observational data available have been taken into account for calculating the (area-weighted) averages.

Figure 17 shows the climatological vertical profiles of the ozone mixing ratio for different latitude bands and months. Some models simulate ozone only up to 10 hPa, which is just below the layer of maximum ozone concentrations in the stratosphere. Although most models capture the trend and magnitude of total column ozone in Antarctica well, the spread of ozone at 10 hPa in the CMIP5 models is quite large for the same region (80°S).



Figure 17. Vertical ozone profile climatologies (2007-2008) at a) 80°N in March, b) the equator in March, and c) at 80°S in October from individual CMIP5 models with interactive chemistry (colored lines) and the ESA CCI ozone data set (solid black line). The multi-model mean (MMM) is shown as a red solid line with one standard deviation of the inter-model spread shown as the light-blue shaded area. For

comparison, also balloon measurements from the Binary Data Base of Profiles (BDBP; Hassler et al., 2008, 2009) are shown for the respective latitudes (2006-2007).

971 Figure 18 shows the zonally averaged climatological seasonal cycle of total column ozone for the 972 CMIP5 multi-model mean, the two satellite-based reference data sets ESA CCI (Figure 18, upper 973 row) and NIWA (Figure 18, lower row), and the differences of the multi-model mean and the 974 two reference data sets. All data sets (models and observations) have been interpolated linearly 975 to the grid of the observations and all grid cells with no observational data have been excluded 976 from the model data sets. The seasonal cycle is calculated from monthly means averaged over 977 the years 1997 to 2010. As expected, the zonal mean seasonal cycle of total column ozone does 978 not differ much between ESA CCI and NIWA for the above mentioned reason. Only the 979 magnitude of ozone is a few DU higher in northern winter in the ESA CCI data set, which can 980 probably also be attributed to the different merging algorithms used to produce the two data sets. 981 The CMIP5 multi-model mean is able to capture the phase and amplitude of total column ozone 982 but tends to slightly overestimate ozone at the equator throughout the year and underestimate 983 total ozone in Antarctica during summer (November through January). The occurrence of very 984 low ozone values in CMIP5 multi-model mean is delayed by about 1 month compared with the 985 observational data sets and lasts a few weeks longer than shown by the observations.



986

Figure 18. Total column ozone climatologies (1997-2010) for (upper row, from left to right) the multimodel mean of CMIP5 models with interactive chemistry (see Table 1), the ESA CCI ozone data set, and the differences between the CMIP5 multi-model mean and the ESA CCI ozone data. The lower row shows the same plots but for the NIWA combined total column ozone data. The model data have been interpolated to the same grid as the observations. In order to calculate the (area-weighted) global annual averages shown above the individual plots, grid cells in the time series without valid observational data have not been taken into account.

The ESA CCI ozone data sets combine all currently available backscatter nadir spectral UV-Vis sensors, i.e. GOME, SCIAMACHY, GOME-2 and OMI (Lerot et al., 2014) resulting in a harmonized product suitable for analyses of long-term ozone trends (WMO, 2014). The reprocessed ozone profiles from 20 years of observations by GOME, SCIAMACHY and
GOME-2 result in a data set of unprecedented accuracy and consistency (Miles et al., 2015;
Keppens et al., 2015) well suited for the evaluation of global coupled climate models with
interactive chemistry.

994 5.8 Greenhouse Gases: XCO₂

995 In order to compare the ESA CCI XCO₂ data set with CMIP5 simulations, only the emission 996 driven simulations (esmHistorical) are used. These simulations were extended until 2014 with 997 results from simulations of the RCP8.5 (esmrcp85). The differences in modeled CO₂ 998 concentrations in the year 2014 between the different emission scenarios (RCP2.6, RCP4.5, 999 RCP8.5) are rather negligible and are therefore not further discussed in the analysis presented 1000 here. Here, we focus on those models of the CMIP5 ensemble that provide all necessary data to 1001 compare with the ESA CCI GHG data for the full time period (2003-2014): BNU-ESM, 1002 CanESM2, CESM1-BGC, FIO-ESM, GFDL-ESM2G, GFDL-ESM2M, MIROC-ESM, MPI-1003 ESM-LR, MRI-ESM1, and NorESM1-ME (see Table 1). These models include an interactive 1004 carbon cycle and performed emission driven simulations in which the emissions rather than the 1005 concentrations of the greenhouse gases are prescribed (Taylor et al., 2012). This allows the 1006 carbon cycle in the models to react to changes in climate by adjusting their carbon fluxes to the 1007 new climate conditions and providing the atmospheric CO₂ concentration as an output 1008 (Friedlingstein et al., 2006).

For comparison of model and satellite data shown in Figure 19 and Figure 20, the model data were interpolated to the grid of the ESA CCI data set $(5^{\circ}x5^{\circ})$ using local area averaging. Grid cells with missing values in the satellite data were also flagged as missing in the model fields. An important characteristic of the ESA CCI data set is that between 2003-2008 measurements are 1013 only over land whereas from 2009-2014 the record contains measurements over land and ocean.1014 The models have been sampled accordingly.

1015 Figure 19 shows the monthly mean time series of XCO₂ comparing ESA CCI data with CMIP5 1016 simulations in four different latitude bands. For all four latitude bands two main features of the 1017 time series are very prominent: firstly, the increase in XCO₂ between 2003 and 2014. The 1018 increase of about 2 ppm per year is consistent with other observations (Ciais, 2013; Jones and 1019 Cox, 2005; Tans and Keeling, 2015) although the absolute values are not directly comparable 1020 since the ESA CCI product is an average of the total atmospheric column of atmospheric CO₂ 1021 with the concentration at higher altitudes increasing more slowly than at the surface due to 1022 mixing (Shia et al., 2006). The CMIP5 multi-model mean shows a positive bias compared with 1023 the ESA CCI data of about 5-10 ppm in all four domains. Particularly the CESM-BGC and the 1024 GFDL-ESM2M models simulate an XCO₂ bias of about two times higher than the bias of the 1025 multi-model mean. The MRI-ESM1 model has the largest negative bias of the models analyzed 1026 here with a bias of about -20 ppm. This agrees with findings by Friedlingstein et al. (2014) and 1027 Hoffman et al. (2014), who analyzed CO₂ simulated by ESMs. Secondly, the seasonal variation 1028 of XCO₂ is more pronounced in the northern hemisphere (30°N-60°N) because of more 1029 vegetation exchanging carbon with the atmosphere. We note again that no ESA CCI XCO_2 data 1030 over the ocean are available before 2009 (see also section 2.8). Since the main anthropogenic 1031 sources of CO₂ are located over land, the CO₂ concentrations over the oceans are slightly lower 1032 than over land. Thus, there is a small discontinuity in the XCO₂ time series shown in Figure 19 in 1033 the beginning of the year 2009 when measurements over the ocean become available and are 1034 included in the calculation of the averages over the different latitude bands. As a consequence of this artifact, the amplitudes of the seasonal cycle in Figure 19 appear slightly reduced in thebeginning of 2009.

1037 The emission driven CMIP5 models simulate a large spread in XCO_2 at all latitude bands mainly 1038 falling outside the observational (1-sigma) uncertainty of the ESA CCI data. The MPI-ESM-LR 1039 model is in good agreement with the annual average XCO_2 values but overestimates the 1040 amplitude of the seasonal cycle compared with the ESA CCI data.



Figure 19. Time series of column averaged carbon dioxide (XCO₂) from 2003 to 2014 from the CMIP5 emission driven simulations for the historical period (2003 to 2005) extended with RCP8.5 simulations (from 2006 to 2014) in comparison with the ESA CCI GHG XCO₂ data. The CMIP5 models are interpolated to the 5°x5° grid of the observations omitting grid cells with no observations. From top left to bottom right: global average, 30°N-60°N, 30°S-30°N, and 60°S-30°S.

- 1042 The spatial distribution of XCO₂ from the CMIP5 models and the ESA CCI data set is compared
- 1043 by analyzing the deviations from the climatological annual averages (2003-2008 and 2009-2014)
- 1044 shown in Figure 20. Because of the trend in XCO₂, we show the two time periods separately to

1045 reduce artifacts caused by XCO_2 data over the ocean only being available in the second half of 1046 the ESA CCI record (2009-2014) (Buchwitz et al., 2015). The CMIP5 models have been 1047 sampled accordingly averaging only over grid cells with observational data available. Over the 1048 continents the ESA CCI data reveal many expected regional features, such as lower XCO₂ 1049 concentrations over the tropical rain forests and the boreal forests in the northern high latitudes 1050 (Buchwitz et al., 2015). This spatial distribution can be expected because in forest regions and 1051 areas with high vegetation more carbon from the atmosphere is taken up by plants via 1052 photosynthesis (Keeling et al., 1995). Higher than global average values are found particularly in 1053 the northern hemisphere over the United States, Europe, Middle East, India, and China. These 1054 basic features are reproduced by the CMIP5 multi-model mean but the annual average XCO_2 1055 values are overestimated by about 6-10 ppm by the models compared with the ESA CCI data in 1056 the time period 2003-2008. This bias in the CMIP5 multi-model mean is found to increase 1057 slightly to 8-12 ppm in the second half of the ESA CCI XCO₂ record (2009-2014), which could 1058 point to possibly slightly too weak carbon sinks in some models (Friedlingstein et al., 2014).



Figure 20. Annual mean XCO₂ climatologies averaged over the years 2003-2008 (top row) and over the years 2009-2014 (bottom row). Shown are deviations from the global annual mean (printed in the right above each panel) for (left) the CMIP5 multi-model mean and (middle) ESA CCI XCO₂. The right panels show the absolute differences between the CMIP5 multi-model mean and ESA CCI XCO₂ data. The CMIP5 results shown are from emission driven historical simulations extended with the respective RCP8.5 scenario.

1060 **6** Summary and outlook

1061 Diagnostics for a subset of the ESA CCI Phase 2 data including the CCIs sea surface 1062 temperature, sea ice, cloud, soil moisture, land cover, aerosol, ozone, and greenhouse gases have 1063 been implemented into the community diagnostics and performance metrics tool ESMValTool. 1064 This enhanced version of the ESMValTool has been applied to evaluate a suite of CMIP5 models 1065 with the new ESA CCI data sets as well as to compare the new data sets with observations that 1066 have already been widely used for model evaluation. The usage of the ESA CCI data in model 1067 evaluation has been demonstrated in overview statistics of the models' global average 1068 performance using RMSD from the climatological mean seasonal cycle as a metric. The ESA 1069 CCI data sets allow for evaluation of new ECVs such as global soil moisture and AOD from fine 1070 particles from global coupled (free running) climate models for which consistent and long-term 1071 observational data sets have not been previously available. For other variables such as total cloud 1072 cover, sea surface temperature, or total ozone columns, the ESA CCI data sets provide the 1073 possibility to compare previously available observational data sets in addition to the models. This 1074 can help to estimate the uncertainty inherent to model evaluations caused by the choice of a 1075 specific observational reference data set for comparison. The error estimates provided as part of 1076 the ESA CCI data sets on a per grid basis help to further assess and quantify what a climate

1077 model can be realistically expected to reproduce. A new extended version of the Taylor diagram 1078 has been presented that includes observational uncertainty estimates and allows to quickly 1079 identify models with a RMSE compared to the observations of less than the observational 1080 uncertainty (also given as RMSE) by simply gauging the figure. The models cannot be expected 1081 to agree perfectly with the observations given the observational uncertainty. In particular for 1082 ECVs with large observational uncertainties such as certain cloud properties this helps to avoid 1083 over-interpreting model biases that cannot be assessed quantitatively and that might depend 1084 significantly on the choice of the reference data set.

1085 In most cases, the ESA CCI data compare well with existing data sets such as, for instance, 1086 MODIS AOD, NIWA total ozone, or NSIDC sea ice concentration. The additional value of 1087 implementing the ESA CCI data sets into the ESMValTool for these quantities lies particularly 1088 in the harmonized and consistently processed data from different platforms and instruments. 1089 Such data can now be used by the climate modeling community to evaluate long-term trends and 1090 variability of selected modeled ECVs. This is particularly relevant to assessing modeled changes 1091 in ECVs related to projected climate change and an important contribution reducing the 1092 uncertainties in the projected climate change scenarios.

The ESMs participating in CMIP6 will be more complex than the models of the CMIP5 generation and include new or more detailed processes such as more sophisticated dynamical vegetation models, sea ice treatment or interactive chemistry and carbon cycle. Future releases of the ESMValTool will therefore not only include further ESA CCIs such as ocean color, sea level, ice sheets and fire, but also additional ECVs from already implemented CCIs such as column averaged methane or additional cloud properties such as, for instance, cloud water path, spectral cloud albedo and cloud optical properties. 1100 The aim is to apply the enhanced version of the ESMValTool presented in this paper for routine 1101 evaluation of ESMs with observations including the ESA CCI data sets within CMIP6. The 1102 CMIP6 results can be analyzed and evaluated together with other evaluation tools and metrics 1103 packages such as PMP as soon as the results become available on the ESGF. The application of 1104 different analysis/evaluation tools in combination with different and independent observational 1105 data sets will help to get a more complete picture of the performance of the quite complex state-1106 of-the-art ESMs, particularly across different ESM domains. This is an important step to identify 1107 domains and processes that would particularly benefit from further model improvements and one 1108 step further to the ultimate goal of improving our understanding of the climate system and 1109 reducing the uncertainties in projections of future climate change.

1110 **Code Availability**

1111 The enhanced version of the ESMValTool presented in this paper is released under the Apache 1112 License, VERSION 2.0. The newly added ESMValTool namelist 'namelist lauer16rse.xml' 1113 includes the diagnostics that can be used to reproduce the figures of this paper. This enhanced 1114 version will be available from the ESMValTool webpage at http://www.esmvaltool.org/ and 1115 from github (https://github.com/ESMValTool-Core/ESMValTool). Users who apply the software 1116 resulting in presentations or papers are kindly asked to cite the ESMValTool documentation 1117 paper (Evring et al., 2016b) alongside with the software doi (doi: 10.17874/ac8548f0315) and 1118 version number. The wider climate community is encouraged to contribute to this effort and to 1119 join the ESMValTool development team for contribution of additional more in-depth diagnostics 1120 for ESM evaluation.

Acknowledgments. Financial support for the development of the enhanced version of the
ESMValTool presented in this study was provided by ESA's Climate Change Initiative Climate
- 1123 Modelling User Group (CMUG). Additionally, funding was received from the European Union's
- 1124 Horizon 2020 research and innovation programme, under grant agreement No. 641816
- 1125 (CRESCENDO). We acknowledge the World Climate Research Programme's Working Group on
- 1126 Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups
- 1127 (listed in Table 1 of this paper) for producing and making their model output available.

1128 **References**

- Adler, R. F., G. J. Huffman, A. Chang, R. Ferraro, P.-P. Xie, J. Janowiak, B. Rudolf, U.
 Schneider, S. Curtis, D. Bolvin, A. Gruber, J. Susskind, P. Arkin, and E. Nelkin (2003), The
 Version-2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis
 (1979-Present), J. Hydrometeorol., 4, 1147-1167.
- Albergel, C., P. de Rosnay, C. Gruhier, J. Muñoz-Sabater, S. Hasenauer, L. Isaksen, Y. Kerr, and
 W. Wagner (2012), Evaluation of remotely sensed and modelled soil moisture products using
 global ground-based in situ observations, Remote Sensing Envir., 118, 215-226.
- Arora, V. K., J. F. Scinocca, G. J. Boer, J. R. Christian, K. L. Denman, G. M. Flato, V. V.
 Kharin, W. G. Lee, and W. J. Merryfield (2011), Carbon emission limits required to satisfy
 future representative concentration pathways of greenhouse gases, Geophys. Res. Lett., 38,
 L05805, doi: 10.1029/2010GL046270.
- Bentsen, M., I. Bethke, J. B. Debernard, T. Iversen, A. Kirkevåg, Ø. Seland, H. Drange, C.
 Roelandt, I. A. Seierstad, C. Hoose, and J. E. Kristjánsson (2013), The Norwegian Earth
 System Model, NorESM1-M Part 1: Description and basic evaluation of the physical
 climate, Geosci. Model Dev., 6, 687-720, doi: 10.5194/gmd-6-687-2013.
- Bi, D., M. Dix, S. Marsland, S. O'Farrell, H. Rashid, P. Uotila, A. Hirst, E. Kowalczyk, M.
 Golebiewski, A. Sullivan, H. Yan, N. Hannah, C. Franklin, Z. Sun, P. Vohralik, I. Watterson,
 X. Zhou, R. Fiedler, M. Collier, Y. Ma, J. Noonan, L. Stevens, P. Uhe, H. Zhu, S. Griffies, R.
 Hill, C. Harris, and K. Puri (2013), The ACCESS coupled model: description, control climate
 and evaluation, Aust. Met. Oceanogr. J., 63, 41-64.
- Bodas-Salcedo, A., M. J. Webb, S. Bony, H. Chepfer, J.-L. Dufresne, S. A. Klein, Y. Zhang, R.
 Marchand, J. M. Haynes, R. Pincus, and V. O. John (2011), COSP Satellite simulation
 software for model assessment, Bull. Amer. Meteorol. Soc., 92, 1023-1043, doi:
 10.1175/2011BAMS2856.1.
- Bodeker, G. E., H. Shiona, and H. Eskes (2005), Indicators of Antarctic Ozone Depletion,
 Atmos. Chem. Phys., 5, 2603-2615.

- Bojinski, S., M. Verstraete, T. C. Peterson, C. Richter, A. Simmons, and M. Zemp (2014), The
 Concept of Essential Climate Variables in Support of Climate Research, Applications, and
 Policy, Bull. Amer. Meteor. Soc., 95, 1431-1443, doi: 10.1175/BAMS-D-13-00047.1.
- Bontemps, S., M. Herold, L. Kooistra, A. van Groenestijn, A. Hartley, O. Arino, I. Moreau, and
 P. Defourny (2012), Revisiting land cover observation to address the needs of the climate
 modeling community, Biogeosciences, 9, 2145-2157, doi: 10.5194/bg-9-2145-2012.
- Bovensmann, H., J. P. Burrows, M. Buchwitz, J. Frerick, S. Noël, V. V. Rozanov, K. V. Chance,
 and A. P. H. Goede (1999), SCIAMACHY: Mission Objectives and Measurement Modes, J.
 Atmos. Sci., 56, 127-150.
- Bräu, M. (2013), Sea-ice in decadal and long-term simulations with the Max Planck Institute
 Earth System Model, Bachelor thesis, Ludwig Maximilian University, Munich, Germany.
- Brovkin, V., T. Raddatz, C. H. Reick, M. Claussen, and V. Gayler (2009), Global biogeophysical
 interactions between forest and climate, Geophys. Res. Lett., 36, L07405, doi:
 10.1029/2009GL037543.
- Brovkin, V., L. Boysen, T. Raddatz, V. Gayler, A. Loew, and M. Claussen (2013), Evaluation of
 vegetation cover and land-surface albedo in MPI-ESM CMIP5 simulations, J. Adv. Model.
 Earth Syst., 5, 48-57, doi: 10.1029/2012MS000169.
- Buchwitz, M., R. de Beek, J. P. Burrows, H. Bovensmann, T. Warneke, J. Notholt, J. F. Meirink,
 A. P. H. Goede, P. Bergamaschi, S. Korner, M. Heimann, and A. Schulz (2005), Atmospheric
 methane and carbon dioxide from SCIAMACHY satellite data: initial comparison with
 chemistry and transport models, Atmos. Chem. Phys., 5, 941-962.
- 1176 Buchwitz, M., M. Reuter, O. Schneising, H. Boesch, I. Aben, M. Alexe, R. Armante, P. 1177 Bergamaschi, H. Bovensmann, D. Brunner, B. Buchmann, J. P. Burrows, A. Butz, F. 1178 Chevallier, A. Chédin, C. D. Crevoisier, S. Gonzi, M. De Mazière, E. De Wachter, R. 1179 Detmers, B. Dils, C. Frankenberg, P. Hahne, O. P. Hasekamp, W. Hewson, J. Hevmann, S. 1180 Houweling, M. Hilker, T. Kaminski, G. Kuhlmann, A. Laeng, T. T. v. Leeuwen, G. 1181 Lichtenberg, J. Marshall, S. Noël, J. Notholt, P. Palmer, R. Parker, M. Scholze, G. P. Stiller, T. Warneke, and C. Zehner (2015), The greenhouse gas project of ESA's climate change 1182 1183 initiative (GHG-CCI): overview, achievements and future plans, Int. Arch. Photogramm, 1184 Remote Sens. Spatial Inf. Sci., XL-7/W3, 165-172.
- 1185Buchwitz, M., and M. Reuter (2016), Merged SCIAMACHY/ENVISAT and TANSO-1186FTS/GOSAT atmospheric column-average dry-air mole fraction of CO2 (XCO2) -1187XCO2_CRDP3_001, Technical Document, http://esgf-1188data1.ceda.ac.uk/thredds/fileServer/esg_obs4mips/esacci/ghg/data/obs4mips/crdp_3/CO2/v1011890/TechNote_SCIAGOSAT_L3_CRDP3_001_XCO2_FINAL.pdf (last accessed 22 September11902016).

- Burrows, J. P., E. Holzle, A. P. H. Goede, H. Visser, and W. Fricke (1995), SCIAMACHY Scanning Imaging Absorption Spectrometer for Atmospheric Chartography, Acta Astronaut,
 35, 445-451.
- Cavalieri, D.J., C. L. Parkinson, P. Gloersen, and H. Zwally (1996), Sea Ice Concentrations from
 Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, Arctic, full record,
 doi: http://dx.doi.org/10.5067/8GQ8LZQVL0VL.
- Cho, K., N. Sasaki, H. Shimoda, T. Sakata, and F. Nishio (1996), Evaluation and Improvement
 of SSM/I Sea Ice Concentration Algorithms for the Sea of Okhotsk, Journal Remote Sensing
 of Japan, 16(2), 47-58.
- Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J.
 Galloway, M. Heimann, C. Jones, C. Le Quéré, R. B. Myneni, S. Piao, and P. Thornton
 (2013), Carbon and Other Biogeochemical Cycles. In: Climate Change 2013: The Physical
 Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the
 Intergovernmental Panel on Climate Change, Stocker, T. F., D. Qin, G.-K. Plattner, M.
 Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Midgley (Ed.),
 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Cionni, I., V. Eyring, J. F. Lamarque, W. J. Randel, D. S. Stevenson, F. Wu, G. E. Bodeker, T.
 G. Shepherd, D. T. Shindell, and D. W. Waugh (2011), Ozone database in support of CMIP5
 simulations: results and corresponding radiative forcing, Atmos. Chem. Phys., 11, 1126711292, doi: 10.5194/acp-11-11267-2011.
- 1211 Clarke, L. E., J. A. Edmonds, H. D. Jacoby, H. Pitcher, J. M. Reilly, and R. Richels (2007),
 1212 Scenarios of greenhouse gas emissions and atmospheric concentrations, Sub-report 2.1a of
 1213 Synthesis and Assessment Product 2.1, Climate Change Science Program and the
 1214 Subcommittee on Global Change Research, Washington DC.
- Coldewey-Egbers, M., D. G. Loyola, M. Koukouli, D. Balis, J.-C. Lambert, T. Verhoelst, J.
 Granville, M. van Roozendael, C. Lerot, R. Spurr, S. M. Frith, and C. Zehner (2015), The
 GOME-type Total Ozone Essential Climate Variable (GTO-ECV) data record from the ESA
 Climate Change Initiative, Atmos. Meas. Tech., 8, 3923-3940, doi: 10.5194/amt-8-39232015.
- Collins, M., S. F. B Tett, and C. Cooper (2001), The Internal Climate Variability of HadCM3, a
 Version of the Hadley Centre Coupled Model without Flux Adjustments, Clim. Dyn., 17 (1),
 61-81.
- Collins W. J., N. Bellouin, M. Doutriaux-Boucher, N. Gedney, T. Hinton, C. D. Jones, S.
 Liddicoat, G. Martin, F. O'Connor, J. Rae, C. Senior, I. Totterdell, S. Woodward, T. Reichler,
 and J. Kim (2008), Evaluation of the HadGEM2 model, Meteorological Office Hadley Centre,
 Technical Note 74.

Collins, W. J., N. Bellouin, M. Doutriaux-Boucher, N. Gedney, P. Halloran, T. Hinton, J. Hughes, C. D. Jones, M. Joshi, S. Liddicoat, G. Martin, F. O'Connor, J. Rae, J., C. Senior, S. Sitch, I. Totterdell, A. Wiltshire, and S. Woodward (2011), Development and evaluation of an Earth-System model - HadGEM2, Geosci. Model Dev., 4, 1051-1075, doi: 10.5194/gmd-4-1051-2011.

- Comiso, J. C., and F. Nishio (2008), Trends in the Sea Ice Cover Using Enhanced and
 Compatible AMSR-E, SSM/I, and SMMR Data, J. Geophys. Res., 113, C02S07, doi:
 10.1029/2007JC0043257.
- Comiso, J. C. (2000), Variability and Trends in Antarctic Surface Temperatures from In Situ and
 Satellite Infrared Measurements, J. Climate, 13, 1674-1696, doi: 10.1175/15200442(2000)013<1674:VATIAS>2.0.CO;2.
- Comiso, J. C. (1995), SSM/I Concentrations Using the Bootstrap Algorithm, NASA Reference
 Publication 1380, 40 pages.
- Crow, W. T., A. A. Berg, M. H. Cosh, A. Loew, B. P. Mohanty, R. Panciera, P. de Rosnay, D.
 Ryu, and J. P. Walker (2012), Upscaling sparse ground-based soil moisture observations for
 the validation of coarse-resolution satellite soil moisture products, Rev. Geophys., 50,
 RG2002, doi: 10.1029/2011RG000372.
- 1244 Curry, J. A., J. L. Schramm, and E. E. Ebert (1995), Sea-Ice Albedo Climate Feedback 1245 Mechanism, J. Climate, 8, 240-247.
- de Leeuw, G., T. Holzer-Popp, S. Bevan, W. Davies, J. Descloitres, R. G. Grainger, J.
 Griesfeller, A. Heckel, S. Kinne, L. Klüser, P. Kolmonen, P. Litvinov, D. Martynenko, P. J.
 R. North, B. Ovigneur, N. Pascal, C. Poulsen, D. Ramon, M. Schulz, R. Siddans, L.
 Sogacheva, D. Tanré, G. E. Thomas, T. H. Virtanen, W. von Hoyningen Huene, M. Vountas,
 and S. Pinnock (2015), Evaluation of seven European aerosol optical depth retrieval
 algorithms for climate analysis, Remote Sensing of Environment, 162, 295-315, doi:
 10.1016/j.rse.04.023.
- Dee, D. P., S. M. Uppala, A. J. Simmons, P. Berrisford, P. Poli, S. Kobayashi, U. Andrae, M. A.
 Balmaseda, G. Balsamo, P. Bauer, P. Bechtold, A. C. M. Beljaars, L. van de Berg, J. Bidlot,
 N. Bormann, C. Delsol, R. Dragani, M. Fuentes, A. J. Geer, L. Haimberger, S. B. Healy, H.
 Hersbach, E. V. Holm, L. Isaksen, P. Kallberg, M. Kohler, M. Matricardi, A. P. McNally, B.
 M. Monge-Sanz, J. J. Morcrette, B. K. Park, C. Peubey, P. de Rosnay, C. Tavolato, J. N.
 Thepaut, F. and Vitart (2011), The ERA-Interim reanalysis: configuration and performance of
 the data assimilation system, Q. J. Roy. Meteor. Soc., 137, 553-597.
- Defourny, P., et al. (2015), ESA Land Cover Climate Change Initiative (ESA LC_cci) data:
 ESACCI-LC-L4-LCCS-Map-300m-P5Y-[2000,2005,2010]-v1.6.1 via Centre for
 Environmental Data Analysis.

- Di Gregorio A. (2005), Land cover classification system: Classification concepts and user
 manual for software version 2; Food & Agriculture Organization of the United Nations, R.,
 Ed.; Italy, p. 190.
- Dils, B., M. Buchwitz, M. Reuter, O. Schneising, H. Boesch, R. Parker, S. Guerlet, I. Aben, T.
 Blumenstock, J. P. Burrows, A. Butz, N. M. Deutscher, C. Frankenberg, F. Hase, O. P.
 Hasekamp, J. Heymann, M. De Maziere, J. Notholt, R. Sussmann, T. Warneke, D. Griffith, V.
 Sherlock, and D. Wunch (2014), The Greenhouse Gas Climate Change Initiative (GHG-CCI):
 comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT
 CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, Atmos
 Meas. Tech., 7, 1723-1744.
- Donlon, C. J., M. Martin, J. Starka, J. Roberts-Jones, E. Fiedler, and W. Wimmer (2012), The
 Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, Remote Sensing
 of Environment, 116, 140-158.
- Donner, L. J., and Coauthors (2011), The Dynamical Core, Physical Parameterizations, and
 Basic Simulation Characteristics of the Atmospheric Component AM3 of the GFDL Global
 Coupled Model CM3, J. Climate, 24, 3483-3519.
- Dorigo, W., A. Gruber, R. De Jeu, W. Wagner, T. Stacke, T., A. Loew, C. Albergel, L. Brocca,
 D. Chung, R. Parinussa, and R. Kidd (2014), Evaluation of the ESA CCI soil moisture
 product using ground-based observations, Remote Sensing of Environment, 162, 380-395.
- Dorigo, W., R. de Jeu, D. Chung, R. Parinussa, Y. Liu, W. Wagner, and D. Fernández-Prieto
 (2012), Evaluating global trends (1988-2010) in harmonized multi-satellite surface soil
 moisture, Geophys. Res. Lett., 39, L18405.
- Dorigo, W. A., W. Wagner, R. Hohensinn, S. Hahn, C. Paulik, A. Xaver, A. Gruber, M. Drusch,
 S. Mecklenburg, P. van Oevelen, A. Robock, and T. Jackson (2011), The international soil
 moisture network: A data hosting facility for global in situ soil moisture measurements,
 Hydrology and Earth System Sciences, 15, 1675-1698, doi: 10.5194/hess-15-1675-2011.
- Dorigo, W. A., A. Xaver, M. Vreugdenhil, A. Gruber, A. Hegyiová, A. D. Sanchis-Dufau, D.
 Zamojski, C. Cordes, W. Wagner, and M. Drusch (2013), Global automated quality control of
 in situ soil moisture data from the international soil moisture network, Vadose Zone Journal,
 doi: 10.2136/vzj2012.0097.
- Dorigo, W.A, D. Chung, A. Gruber, S. Hahn, T. Mistelbauer, R. M. Parinussa, C. Paulik, C.
 Reimer, R. van der Schalie, R. A. M. de Jeu, and W. Wagner (2016), [Hydrological Cycle]
 Soil Moisture [in: "State of the Climate 2015"], Bull. Am. Meteorol. Soc., 97(8), S31-S32.
- Du, E. A. Di Vittorio, W. D. Collins (2016), Evaluation of hydrologic components of community
 land model 4 and bias identification, Int. J. Appl. Earth Observ. Geoinform., 48, 5-16, doi:
 10.1016/j.jag.2015.03.013.

1299 Dufresne, J.-L., M.-A. Foujols, S. Denvil, A. Caubel, O. Marti, O. Aumont, Y. Balkanski, S. 1300 Bekki, H. Bellenger, R. Benshila, S. Bony, L. Bopp, P. Braconnot, P. Brockmann, P. Cadule, F. Cheruy, F. Codron, A. Cozic, D. Cugnet, N. de Noblet, J.-P. Duvel, C. Ethé, L. Fairhead, T. 1301 1302 Fichefet, S. Flavoni, P. Friedlingstein, J.-Y. Grandpeix, L. Guez, E. Guilvardi, D. 1303 Hauglustaine, F. Hourdin, A. Idelkadi, J. Ghattas, S. Joussaume, M. Kageyama, G. Krinner, S. 1304 Labetoulle, A. Lahellec, M.-P. Lefebvre, F. Lefevre, C. Levy, Z. X. Li, J. Lloyd, F. Lott, G. Madec, M. Mancip, M. Marchand, S. Masson, Y. Meurdesoif, J. Mignot, I. Musat, S. Parouty, 1305 1306 J. Polcher, C. Rio, M. Schulz, D. Swingedouw, S. Szopa, C. Talandier, P. Terray, N. Viovy, 1307 N. Vuichard (2013), Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, Clim Dyn, 40, 2123-2165, doi: 10.1007/s00382-012-1636-1. 1308

Embury, O., C. J. Merchant, and M. J. Filipiak (2012a), A reprocessing for climate of sea surface
temperature from the Along-Track Scanning Radiometers: basis in radiative transfer, Remote
Sensing of Environment, 116, 32-46, doi: 10.1016/j.rse.2010.10.016.

Embury, O., C. J. Merchant, and G. K. Corlett (2012b), A reprocessing for climate of sea surface
temperature from the Along-Track Scanning Radiometers: initial validation, accounting for
skin and diurnal variability, Remote Sensing of Environment, 116, 62-78, doi:
10.1016/j.rse.2011.02.028.

- Eyring, V., J. M. Arblaster, I. Cionni, J. Sedlacek, J. Perlwitz, P. J. Young, S. Bekki, D.
 Bergmann, P. Cameron-Smith, W. J. Collins, G. Faluvegi, K.-D. Gottschaldt, L. W. Horowitz,
 D. E. Kinnison, J.-F. Lamarque, D. R. Marsh, D. Saint-Martin, D. T. Shindell, K. Sudo, S.
 Szopa, and S. Watanabe (2013), Long-Term Ozone Changes and Associated Climate Impacts
 in CMIP5 Simulations, J. Geophys. Res., 118(10), 5029-60.
- Eyring, V., S. Bony, G. A. Meehl, C. A. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor
 (2016a), Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)
 experimental design and organization, Geosci. Model Dev., 9, 1937-1958, doi: 10.5194/gmd9-1937-2016.
- Eyring, V., M. Righi, A. Lauer, M. Evaldsson, S. Wenzel, C. Jones, A. Anav, O. Andrews, I.
 Cionni, E. L. Davin, C. Deser, C. Ehbrecht, P. Friedlingstein, P. Gleckler, K.-D. Gottschaldt,
 S. Hagemann, M. Juckes, S. Kindermann, J. Krasting, D. Kunert, R. Levine, A. Loew, J.
 Mäkelä, G. Martin, E. Mason, A. S. Phillips, S. Read, C. Rio, R. Roehrig, D. Senftleben, A.
 Sterl, L. H. van Ulft, J. Walton, S. Wang, and K. D. Williams (2016b), ESMValTool (v1.0) –
 a community diagnostic and performance metrics tool for routine evaluation of Earth System
 Models in CMIP, Geosci. Model Dev., 9, 1747-1802, doi: 10.5194/gmd-9-1747-2016.
- Eyring, V., P. J. Gleckler, C. Heinze, R. J. Stouffer, K. E. Taylor, V. Balaji, E. Guilyardi, S.
 Joussaume, S. Kindermann, B. N. Lawrence, G. A. Meehl, M. Righi, and D. N. Williams
 (2016c), Towards improved and more routine Earth system model evaluation in CMIP, Earth
 Syst. Dynam., 7, 813-830, doi: 10.5194/esd-7-813-2016.

- Ferraro, R., D. E. Waliser, P. Gleckler, K. E. Taylor, and V. Eyring (2015), Evolving obs4MIPs
 to Support the Sixth Coupled Model Intercomparison Project (CMIP6), B. Am. Meteorol.
 Soc., doi: 10.1175/BAMS-D-14-00216.1.
- Flato, G., J. Marotzke, B. Abiodun, P. Braconnot, S. C. Chou, W. Collins, P. Cox, F. Driouech,
 S. Emori, V. Eyring, C. Forest, P. Gleckler, E. Guilyardi, C. Jakob, V. Kattsov, C. Reason,
 and M. Rummukainen (2013), Evaluation of Climate Models. In: Climate Change 2013: The
 Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of
 the Intergovernmental Panel on Climate Change, Stocker, T. F., D. Qin, G.-K. Plattner, M.
- 1344 Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley (Ed.), 1345 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Friedlingstein, P., P. Cox, R. Betts, L. Bopp, W. von Bloh, V. Brovkin, P. Cadule, S. Doney, M.
 Eby, I. Fung, G. Bala, J. John, C. Jones, F. Joos, T. Kato, M. Kawamiya, W. Knorr, K.
 Lindsay, H. D. Matthews, T. Raddatz, P. Rayner, C. Reick, E. Roeckner, K.-G. Schnitzler, R.
 Schnur, K. Strassmann, A. J. Weaver, C. Yoshikawa, and N. Zeng (2006), Climate-Carbon
 Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison, J. Climate, 19,
 3337-3353.
- Friedlingstein, P., M. Meinshausen, V. K. Arora, C. D. Jones, A. Anav, S. K. Liddicoat, and R.
 Knutti (2014), Uncertainties in CMIP5 Climate Projections due to Carbon Cycle Feedbacks, J.
 Climate, 27, 511-526, doi: 10.1175/JCLI-D-12-00579.1.
- Gent, P. R., G. Danabasoglu, L. J. Donner, M. M. Holland, E. C. Hunke, S. Jayne, D. M.
 Lawrence, R. B. Neale, P. J. Rasch, M. Vertenstein, P. H. Worley, Z.-L. Yang, and M. Zhang
 (2011), The Community Climate System Model version 4, J. Climate, 24(19), 4973-4991, doi:
 10.1175/2011JCLI4083.1.
- GCOS (2010), Implementation Plan for the Global Observing System for Climate in Support ofthe UNFCCC, August 2010.
- Gleckler, P. J., K. E. Taylor, and C. Doutriaux (2008), Performance metrics for climate models,
 J. Geophys. Res., 113, D06104.
- Gleckler, P. J., C. Doutriaux, P. J. Durack, K. E. Taylor, Y. Zhang, D. N. Williams, E. Mason,
 and J. Servonnat (2016), A More Powerful Reality Test for Climate Models, Eos Trans. AGU,
 doi: 10.1029/2016EO051663.
- Good, S, and N. Rayner (2014), Product Specification Document, Project Document SST_CCI PSD-UKMO-201, http://www.esa-sst-cci.org/PUG/documents (last accessed 13 July 2016).
- Hagemann, S., A. Loew, and A. Andersson (2013), Combined evaluation of MPI-ESM land
 surface water and energy fluxes, J. Adv. Model. Earth Syst., 5, 259-286, doi:
 10.1029/2012MS000173.

- Hassler, B., G. E. Bodeker, and M. Dameris (2008), Technical Note: A new global database of
 trace gases and aerosols from multiple sources of high vertical resolution measurements,
 Atmos. Chem. Phys., 8, 5403-5421.
- Hassler, B., G. E. Bodeker, I. Cionni, and M. Dameris (2009), A vertically resolved, monthly
 mean, ozone database from 1979 to 2100 for constraining global climate model simulations,
 International Journal of Remote Sensing, 30(15-16), 4009-4018.
- Hazeleger, W., C. Severijns, T. Semmler, S. Ştefănescu, S. Yang, X. Wang, K. Wyser, E. Dutra,
 J. M. Baldasano, R. Bintanja, P. Bougeault, R. Caballero, A. M. L. Ekman, J. H. Christensen,
 B. van den Hurk, P. Jimenez, C. Jones, P.Kållberg, T. Koenigk, R. Mc Grath, P. Miranda, T.
 van Noije, T. Palmer, J. A. Parodi, T. Schmith, F. Selten, T. Storelvmo, A. Sterl, H. Tapamo,
 M. Vancoppenolle, P. Viterbo, and U. Willén (2010), EC-Earth, A Seamless Earth-System
 Prediction Approach in Action, Bull. Am. Meteor. Soc., 91, 1357-1363, doi:
 10.1175/2010BAMS2877.1.
- Heidinger, A. K., M. J. Foster, A. Walther, and X. Zhao (2014), The Pathfinder atmospheresextended AVHRR climate dataset, Bull. Am. Meteor. Soc., 95(6), 909-922, doi:
 10.1175/BAMS-D-12-00246.1.
- Hoffman, F. M., J. T. Randerson, V. K. Arora, Q. Bao, P. Cadule, D. Ji, C. D. Jones, M. Kawamiya, S. Khatiwala, K. Lindsay, A. Obata, E. Shevliakova, K. D. Six, J. F. Tjiputra, E. M. Volodin, and T. Wu (2014), Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models, J. Geophys. Res. Biogeosci., 119, 141-162, doi: 10.1002/2013JG002381.
- Holben, B. N., T. F. Eck, I. Slutsker, D. Tanré, J. P. Buis, A. Setzer, E. Vermote, J. A. Reagan,
 Y. J. Kaufman, T. Nakajima, F. Lavenu, I. Jankowiak, and A. Smirnov (1998), AERONET –
 A Federated Instrument Network and Data Archive for Aerosol Characterization, Remote
 Sens. Environ., 66, 1-16, doi: 10.1016/s0034-4257(98)00031-5.
- Hollmann, R., C. J. Merchant, R. Saunders, C. Downy, M. Buchwitz, A. Cazenave, W. Wagner
 (2013), The ESA climate change initiative: Satellite data records for essential climate
 variables, Bull. Am. Meteor. Soc., 94(10), 1541-1552, doi: 10.1175/BAMS-D-11-00254.1.
- Hourdin, F., M.-A. Foujols, F. Codron, V. Guemas, J.-L. Dufresne, S. Bony, S. Denvil, L. Guez,
 F. Lott, J. Ghattas, P. Braconnot, O. Marti, Y. Meurdesoif, L. Bopp (2013), Impact of the
 LMDZ atmospheric grid configuration on the climate and sensitivity of the IPSL-CM5A
 coupled model, Clim. Dyn., 40, 2167-2192, doi: 10.1007/s00382-012-1411-3.
- Hubert, D., J.-C. Lambert, T. Verhoelst, J. Granville, A. Keppens, J.-L. Baray, A. E. Bourassa,
 U. Cortesi, D. A. Degenstein, L. Froidevaux, S. Godin-Beekmann, K. W. Hoppel, B. J.
 Johnson, E. Kyrölä, T. Leblanc, G. Lichtenberg, M. Marchand, C. T. McElroy, D. Murtagh,
 H. Nakane, T. Portafaix, R. Querel, J. M. Russell III, J. Salvador, H. G. J. Smit, K. Stebel, W.
 Steinbrecht, K. B. Strawbridge, R. Stübi, D. P. J. Swart, G. Taha, D. W. Tarasick, A. M.

- Thompson, J. Urban, J. A. E. van Gijsel, R. Van Malderen, P. von der Gathen, K. A. Walker,
 E. Wolfram, and J. M. Zawodny (2016), Ground-based assessment of the bias and long-term
 stability of 14 limb and occultation ozone profile data records, Atmos. Meas. Tech., 9, 24972534, doi: 10.5194/amt-9-2497-2016.
- Hung, K.C, H. Gu, and Z. A. Zong (2004), A modified superconvergent patch recovery method
 and its application to large deformation problems, Finite Elements in Analysis and Design,
 40(5-6).
- Huffman, G. J., and D. T. Bolvin (2013), GPCP Version 2.2 SG Combined Precipitation Data
 Set Documentation, available online at ftp://precip.gsfc.nasa.gov/pub/gpcpv2.2/doc/V2.2_doc.pdf (last accessed 13 September 2016), http://precip.gsfc.nasa.gov.
- Hurrell, J. W., M. M. Holland, P. R. Gent, J. E. Kay, P. J. Kushner, J.-F. Lamarque, W. G. Large,
 D. Lawrence, K. Lindsay, W. H. Lipscomb, M. C. Long, N. Mahowald, D. R. Marsh, R. B.
 Neale, P. Rasch, S. Vavrus, M. Vertenstein, D. Bader, W. D. Collins, J. J. Hack, J. Kiehl, and
 S. Marshall (2013), The Community Earth System Model: A Framework for Collaborative
 Research, Bull. Am. Meteor. Soc., 94(9), 1339-1360, doi: 10.1175/BAMS-D-12-00121.1.
- Ivanova, N., L. T. Pedersen, R. T. Tonboe, S. Kern, G. Heygster, T. Lavergne, A. Sørensen, R.
 Saldo, G. Dybkjær, L. Brucker, and M. Shokr (2015), Inter-comparison and evaluation of sea
 ice algorithms: towards further identification of challenges and optimal approach using
 passive microwave observations, The Cryosphere, 9, 1797-1817, doi: 10.5194/tc-9-17972015.
- Ji, D., L. Wang, J. Feng, Q. Wu, H. Cheng, Q. Zhang, J. Yang, W. Dong, Y. Dai, D. Gong, R.-H.
 Zhang, X. Wang, J. Liu, J. C. Moore, D. Chen, and M. Zhou (2014), Description and basic
 evaluation of Beijing Normal University Earth System Model (BNU-ESM) version 1, Geosci.
 Model Dev., 7, 2039-2064, doi: 10.5194/gmd-7-2039-2014.
- Jones, C. D., and P. M. Cox (2005), On the significance of atmospheric CO₂ growth rate
 anomalies in 2002-2003, Geophys. Res. Lett., 32.
- 1434 Kalnay, E., et al. (1996), The NCEP/NCAR 40-Year Reanalysis Project, Bull. Am. Meteor. Soc.,
 1435 77, 437-471, doi: 10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2.
- Karlsson, K.-G., A. Riihelä, R. Müller, J. F. Meirink, J. Sedlar, M. Stengel, M. Lockhoff, J.
 Trentmann, F. Kaspar, R. Hollmann, and E. Wolters (2013), CLARA-A1: a cloud, albedo,
 and radiation dataset from 28 yr of global AVHRR data, Atmos. Chem. Phys., 13, 5351-5367,
 doi: 10.5194/acp-13-5351-2013.
- Keeling, C. D., T. P. Whorf, M. Wahlen, and J. Vanderplicht (1995), Interannual Extremes in the
 Rate of Rise of Atmospheric Carbon-Dioxide since 1980, Nature, 375, 666-670.
- Keppens, A., J.-C. Lambert, J. Granville, G. Miles, R. Siddans, J. C. A. van Peet, R. J. van der A,
 D. Hubert, T. Verhoelst, A. Delcloo, S. Godin-Beekmann, R. Kivi, R. Stübi, and C. Zehner

- (2015), Round-robin evaluation of nadir ozone profile retrievals: methodology and application
 to MetOp-A GOME-2, Atmos. Meas. Tech., 8, 2093-2120, doi:10.5194/amt-8-2093-2015.
- Kern, S., K. Khvorostovsky, H. Skourup, E. Rinne, Z. S. Parsakhoo, V. Djepa, P. Wadhams, and
 S. Sandven (2015), The impact of snow depth, snow density and ice density on sea ice
 thickness retrieval from satellite radar altimetry: results from the ESA-CCI Sea Ice ECV
 Project Round Robin Exercise, The Cryosphere, 9, 37-52, doi: 10.5194/tc-9-37-2015.
- Kern, S., A. Rösel, L. T. Pedersen, N. Ivanova, R. Saldo, and R. T. Tonboe (2016), The impact
 of melt ponds on summertime microwave brightness temperatures and sea ice concentrations,
 The Cryosphere Discuss., doi: 10.5194/tc-2015-202.
- Khoei, S. A., and A. R. Gharehbaghi (2007), The superconvergent patch recovery technique and
 data transfer operators in 3d plasticity problems, Finite Elements in Analysis and Design,
 43(8).
- Koukouli, M., D. Balis, I. Zyrichidou, C. Lerot, M. Van Roozendael, J-C. Lambert, J. Granville,
 J-P. Pommereau, F. Goutail, G. Labow, S. Frith, D. Loyola, R. Spurr, and C. Zehner (2015),
 Evaluating a new homogeneous total ozone climate data record from GOME/ERS-2,
 SCIAMACHY/Envisat, and GOME-2/MetOp-A, J. Geophys. Res., 120(23), 12,296-12,312,
 doi: 10.1002/2015JD023699.
- Kuze, A., H. Suto, M. Nakajima, and T. Hamazaki (2009), Thermal and near infrared sensor for
 carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing
 Satellite for greenhouse gases monitoring, Appl. Opt., 48, 6716-6733.
- Kwok R, and G. F. Cunningham (2015), Variability of Arctic sea ice thickness and volume from
 CryoSat-2, Phil. Trans. R. Soc. A., 373, 20140157, doi: 10.1098/rsta.2014.0157.
- Lauer, A., and K. Hamilton (2013), Simulating Clouds with Global Climate Models: A
 Comparison of CMIP5 Results with CMIP3 and Satellite Data, J. Climate, 26, 3823-3845,
 doi: 10.1175/JCLI-D-12-00451.1.
- Lavergne, T., and E. Rinne (2014), ESA Sea Ice Climate Change Initiative: Phase 1, D3.4
 Product User Guide (PUG), doc ref: SICCI-PUG-13-07, version: 2.0, date: 29 August 2014.
- Lerot, C., M. Van Roozendael, R. Spurr, et al. (2014), Homogenized total ozone data records
 from the European sensors GOME/ERS-2, SCIAMACHY/Envisat and GOME-2/Metop-A, J.
 Geophys. Res., 119, 1-20, doi: 10.1002/2013JD020831.
- Levy, R. C., S. Mattoo, L. A. Munchak, L. A. Remer, A. M. Sayer, F. Patadia, and N. C. Hsu
 (2013), The Collection 6 MODIS aerosol products over land and ocean, Atmos. Meas. Tech.,
 6, 2989-3034, doi: 10.5194/amt-6-2989-2013.
- Li, L., B. Wang, Y. Yu, et al. (2010), Development and evaluation of GAMIL2.0 and
 FGOALS2.0-g. In: The 5th C20C Workshop, 25-28 October 2010, IAP, Beijing.

- Li, W., P. Ciais, N. MacBean, Shushi Penga, P. Defourny, and S. Bontemps (2016), Major forest
 changes and land cover transitions based on plant functional types derived from the ESA CCI
 Land Cover product, Int. J. Appl. Earth Observ. Geoinform., 47, 30-39, doi:
 10.1016/j.jag.2015.12.006.
- Lindsay, R., and A. Schweiger (2015), Arctic sea ice thickness loss determined using subsurface,
 aircraft, and satellite observations, The Cryosphere, 9, 269-283, doi: 10.5194/tc-9-269-2015.
- Liu, Y. Y., R. M. Parinussa, W. A. Dorigo, R. A. M. de Jeu, W. Wagner, A. I. J. M. van Dijk, M.
 F. McCabe, and J. P. Evans (2011), Developing an improved soil moisture dataset by
 blending passive and active microwave satellite-based retrievals, Hydrology and Earth System
 Sciences, 15, 425-436, doi: 10.5194/hess-15-425-2011.
- Liu, Y. Y., W. A. Dorigo, R. M. Parinussa, R. A. M. de Jeu, W. Wagner, M. F. McCabe, J. P.
 Evans, and A. I. J. M. van Dijk (2012), Trend-preserving blending of passive and active
 microwave soil moisture retrievals, Remote Sensing of Environment, 123, 280-297, doi:
 10.1016/j.rse.2012.03.014.
- Loeb, N. G., B. A. Wielicki, D. R. Doelling, G. L. Smith, D. F. Keyes, S. Kato, N. ManaloSmith, T. Wong (2009), Toward optimal closure of the Earth's top-of-atmosphere radiation
 budget, J. Climate, 22, 748-766, doi: 10.1175/2008JCLI2637.1.
- Loeb, N. G., J. M. Lyman, G. C. Johnson, R. P. Allan, D. R. Doelling, T. Wong, B. J. Soden, and
 G. L. Stephens (2012), Observed changes in top-of-the-atmosphere radiation and upper-ocean
 heating consistent within uncertainty, Nat. Geosci., 5, doi: 10.1038/NGEO1375.
- Loew, A., T. Stacke, W. Dorigo, R. de Jeu, and S. Hagemann (2013), Potential and limitations of
 multidecadal satellite soil moisture observations for selected climate model evaluation studies,
 Hydrology and Earth System Sciences, 17, 3523-3542.
- Loew, A. (2008), Impact of surface heterogeneity on surface soil moisture retrievals from
 passive microwave data at the regional scale: The upper danube case, Remote Sensing of
 Environment, 112, 231-248, doi: 10.1016/j.rse.2007.04.009.
- Long, C. M., Lindsay, K., S. Peacock, J. K. Moore, and S. C. Doney (2013), Twentieth-Century
 Oceanic Carbon Uptake and Storage in CESM1(BGC)*, J. climate, 26, 6775-6800, doi:
 10.1175/JCLI-D-12-00184.1.
- Loyola, D. G., R. M. Coldewey-Egbers, M. Dameris, H. Garny, A. Stenke, M. Van Roozendael,
 C. Lerot, D. Balis, and M. Koukouli (2009), Global long-term monitoring of the ozone layer a prerequisite for predictions, International Journal of Remote Sensing, 30, 15, 4295-4318,
 doi: 10.1080/01431160902825016.
- Marsh, D. R., M. J. Mills, D. E. Kinnison, J.-F. Lamarque, N. Calvo, and L. M. Polvani (2013),
 Climate Change from 1850 to 2005 Simulated in CESM1(WACCM), J. Climate, 26, 73727391, doi: 10.1175/JCLI-D-12-00558.1.

- Mcgarragh G., C. Poulsen, G. Thomas, A. Povey, O. Sus, C. Schlundt, S. Stapelberg, S. Proud,
 M. Christensen, M. Stengel, and R. Grainger (2016), The Community Cloud retrieval for
 CLimate (CC4CL). Part II: The optimal estimation algorithm, manuscript in preparation.
- Merchant, C. J., O. Embury, J. Roberts-Jones, E. Fiedler, C. E. Bulgin, G. K. Corlett, S. Good,
 A. McLaren, N. Rayner, S. Morak-Bozzo, and C. Donlon (2014a), Sea surface temperature
 datasets for climate applications from Phase 1 of the European Space Agency Climate Change
 Initiative (SST CCI), Geoscience Data Journal, 1, 179-191, doi: 10.1002/gdj3.20.
- Merchant, C. J., O. Embury, J. Roberts-Jones, E. K. Fiedler, C. E. Bulgin, G. K. Corlett, S.
 Good, A. McLaren, N. A. Rayner, and C. Donlon (2014b), ESA Sea Surface Temperature
 Climate Change Initiative (ESA SST CCI): Analysis long term product version 1.0, NERC
 Earth Observation Data Centre, 24th February 2014, doi: 10.5285/878bef44-d32a-40cd-a02d49b6286f0ea4.
- Miles, G. M., R. Siddans, B. J. Kerridge, B. G. Latter, and N. A. D. Richards (2015),
 Tropospheric ozone and ozone profiles retrieved from GOME-2 and their validation, Atmos.
 Meas. Tech., 8, 385-398, doi: 10.5194/amt-8-385-2015.
- Nam, C., S. Bony, J. L. Dufresne, and H. Chepfer (2012), The 'too few, too bright' tropical lowcloud problem in CMIP5 models, Geophys. Res. Lett., 39, L21801, doi:
 10.1029/2012GL053421.
- Parinussa, R. M., A. G. C. A. Meesters, Y. Y. Liu, W. Dorigo, W. Wagner, and R. A. de Jeu
 (2011), Error estimates for near-real-time satellite soil moisture as derived from the land
 parameter retrieval model, IEEE Geoscience and Remote Sensing Letters, 8, 779-783, doi:
 10.1109/LGRS.2011.2114872.
- Phillips, A. S., C. Deser, and J. Fasullo (2014), Evaluating Modes of Variability in Climate
 Models, Eos Trans. AGU, 95(49), 453-455.
- 1539 Platnick, S., M. D. King, S. A. Ackerman, W. P. Menzel, B. A. Baum, J. C. Riedi, and R. A. 1540 Frey (2003), The MODIS cloud products: algorithms and examples from Terra, IEEE Transactions Geoscience 1541 on and Remote Sensing. 41. 2. 459-473, doi: 1542 10.1109/TGRS.2002.808301.
- Popp, T., G. de Leeuw, C. Bingen, C. Brühl, V. Capelle, A. Chedin, L. Clarisse, O. Dubovik, R.
 Grainger, J. Griesfeller, A. Heckel, S. Kinne, L. Klüser, M. Kosmale, P. Kolmonen, L. Lelli,
 P. Litvinov, L. Mei, P. North, S. Pinnock, A. Povey, C. Robert, M. Schulz, L. Sogacheva, K.
 S., D. S. Zweers, G. Thomas, L. Gijsbert Tilstra, S. Vandenbussche, P. Veefkind, M. Vountas,
 and Y. Xue (2016), Development, Production and Evaluation of Aerosol Climate Data
 Records from European Satellite Observations (Aerosol_cci), Remote Sensing, 8, 421; doi:
 10.3390/rs8050421.
- Poulter, B., N. MacBean, A. Hartley, I. Khlystova, O. Arino, R. Betts, S. Bontemps, M.
 Boettcher, C. Brockmann, P. Defourny, S. Hagemann, M. Herold, G. Kirches, C. Lamarche,

- D. Lederer, C. Ottlé, M. Peters, and P. Peylin (2015), Plant functional type classification for
 earth system models: results from the European Space Agency's Land Cover Climate Change
 Initiative, Geosci. Model Dev., 8, 2315-2328, doi: 10.5194/gmd-8-2315-2015.
- Qiao, F., Z. Song, Y. Bao, Y. Song, Q. Shu, C. Huang, and W. Zhao (2013), Development and
 evaluation of an Earth System Model with surface gravity waves, J. Geophys. Res. Oceans,
 118, 4514-4524, doi: 10.1002/jgrc.20327.
- Qiu, J., Q. Gao, S. Wang, and Z. Su (2016), Comparison of temporal trends from multiple soil
 moisture data sets and precipitation: The implication of irrigation on regional soil moisture
 trend, Int. J. Appl. Earth Observ. Geoinform., 48, 17-27, doi: 10.1016/j.jag.2015.11.012.
- Rayner, N. A., D. E. Parker, E. B. Horton, C. K. Folland, L. V. Alexander, D. P. Rowell, E. C.
 Kent, and A. Kaplan (2003), Global analyses of sea surface temperature, sea ice, and night
 marine air temperature since the late nineteenth century, J. Geophys. Res., 108, 4407.
- Rayner, N, S. Good, and T. Block (2015), SST CCI Product User Guide, Project Document
 SST_CCI-PUG-UKMO-201, http://www.esa-sst-cci.org/PUG/documents, 2015 (last accessed
 13 July 2016).
- Remer, L. A., Y. J. Kaufman, D. Tanré, S. Mattoo, D. A. Chu, J. V. Martins, R. R. Li, C. Ichoku,
 R. C. Levy, R. G. Kleidman, T. F. Eck, E. Vermote, and B. N. Holben (2005), The MODIS
 Aerosol Algorithm, Products, and Validation, J. Atmos. Sci., 62, 947-973, doi:
 10.1175/jas3385.1.
- Reuter, M., M. Buchwitz, O. Schneising, F. Hase, J. Heymann, S. Guerlet, A. J. Cogan, H.
 Bovensmann, and J. P. Burrows (2012), A simple empirical model estimating atmospheric
 CO2 background concentrations, Atmos. Meas. Tech., 5, 1349-1357.
- Reynolds, R. W., D. B. Chelton, J. Roberts-Jones, M. J. Martin, D. Menemenlis, and C. J.
 Merchant (2013), Objective Determination of Feature Resolution in Two Sea Surface
 Temperature Analyses, J. Climate, 26, 2514-2533, doi: 10.1175/JCLI-D-12-00787.1.
- 1577 Ricker, R., S. Hendricks, V. Helm, H. Skourup, and M. Davidson (2014), Sensitivity of CryoSat-
- 1578 2 Arctic sea-ice freeboard and thickness on radar-waveform interpretation, The Cryosphere, 8,
 1579 1607-1622, doi: 10.5194/tc-8-1607-2014.
- Roberts-Jones, J., E. K. Fiedler, and M. J. Martin (2012), Daily, Global, High-Resolution SST
 and Sea Ice Reanalysis for 1985-2007 Using the OSTIA System, J. Climate, 25, 6215-6232,
 doi: 10.1175/JCLI-D-11-00648.1.
- Robert-Jones, J., K. Bovis, M. J. Martin, and A. McLaren (2016), Estimating background error
 covariance parameters and assessing their impact in the OSTIA system, Remote Sensing of
 Environment, 176, 117-138, doi: 10.1016/j.rse.2015.12.006.
- Rodgers, C. D. (2000), Inverse Methods for Atmospheric Sounding: Theory and Practice, World
 Scientific Publishing.

- Roeckner, E., R. Brokopf, M. Esch, M. Giorgetta, S. Hagemann, L. Kornblueh, E. Manzini, U.
 Schlese, and U. Schulzweida (2006), Sensitivity of simulated climate to horizontal and
 vertical resolution in the ECHAM5 atmosphere model, J. Climate, 19, 3771-3791.
- Rotstayn, L., M. Collier, M. Dix, Y. Feng, Y., H. Gordon, S. O'Farrell, I. Smith, J. and Syktus (2010), Improved simulation of Australian climate and ENSO-related climate variability in a GCM with an interactive aerosol treatment, Int. J. Climatology, 30(7), 1067-1088, doi: 10.1002/joc.1952.
- Sakamoto and Coauthors (2012), MIROC4h a new high-resolution atmosphere-ocean coupled
 general circulation model, Journal of the Meteorological Society of Japan, 90(3), 325-359,
 doi: 10.2151/jmsj.2012-301.
- Sandven, S., et al. (2015), ESA Sea Ice Climate Change Initiative (ESA Seaice_cci) data: ESA
 CCI SIC v1.11, via Centre for Environmental Data Analysis, 16 Feb 2016.
- Santoro, M., U. Wegmüller (2014), Multi-temporal Synthetic Aperture Radar Metrics Applied to
 Map Open Water Bodies, IEEE Journal of Selected Topics in Applied Earth Observations and
 Remote Sensing 2014, 7, 3225-3238.
- Schmidt, G. A., and Coauthors (2006), Present day atmospheric simulations using GISS ModelE:
 Comparison to in-situ, satellite and reanalysis data, J. Climate, 19, 153-192.
- Scoccimarro E., S. Gualdi, A. Bellucci, A. Sanna, P.G. Fogli, E. Manzini, M. Vichi, P. Oddo,
 and A. Navarra (2011), Effects of Tropical Cyclones on Ocean Heat Transport in a High
 Resolution Coupled General Circulation Model, J. Climate, 24, 4368-4384.
- Shia, R.-L., M.-C. Liang, C. E. Miller, and Y. L. Yung (2006), CO₂ in the upper troposphere:
 Influence of stratosphere-troposphere exchange, Geophys. Res. Lett., 33, L14814, doi:
 10.1029/2006GL026141.
- Smirnov, A., B. N. Holben, I. Slutsker, D. M. Giles, C. R. McClain, T. F. Eck, S. M. Sakerin, A.
 Macke, P. Croot, G. Zibordi, P. K. Quinn, J. Sciare, S. Kinne, M. Harvey, T. J. Smyth, S.
 Piketh, T. Zielinski, A. Proshutinsky, J. I. Goes, N. B. Nelson, P. Larouche, V. F. Radionov,
 P. Goloub, K. Krishna Moorthy, R. Matarrese, E. J. Robertson, and F. Jourdin (2009),
 Maritime Aerosol Network as a component of Aerosol Robotic Network, J. Geophys. Res.,
 114, D06204, doi: 10.1029/2008JD011257.
- 1617 Smith, S. J., and T. M. L. Wigley (2006), MultiGas forcing stabilization with minicam, The 1618 Energy Journal Special issue, 3, 373-392.
- Sofieva, V. F., N. Rahpoe, J. Tamminen, E. Kyrölä, N. Kalakoski, M. Weber, A. Rozanov, C.
 von Savigny, A. Laeng, T. von Clarmann, G. Stiller, S. Lossow, D. Degenstein, A. Bourassa,
 C. Adams, C. Roth, N. Lloyd, P. Bernath, R. J. Hargreaves, J. Urban, D. Murtagh, A.
 Hauchecorne, F. Dalaudier, M. van Roozendael, N. Kalb, and C. Zehner (2013), Harmonized
 dataset of ozone profiles from satellite limb and occultation measurements, Earth Syst. Sci.
 Data, 5, 349-363, doi: 10.5194/essd-5-349-2013.

- Stengel M., C. Poulsen, G. Thomas, M. Christiansen, C. Carbajal Henken, R. Preusker, J.
 Fischer, O. Sus, S. Stapelberg, C. Schlundt, A. Devasthale, U. Willén, K.-G. Karlsson, A.
 Povey, D. Grainger, S. Proud, J. F. Meirink, A. Feofilov, and R. Hollmann (2016a), Cloud
 property datasets retrieved from AVHRR, MODIS, AATSR and MERIS in the framework of
 the Cloud cci project, manuscript in preparation.
- Stengel M., O. Sus, S. Stapelberg, C. Schlundt, C. Poulsen, and R. Hollmann (2016b), ESA
 Cloud Climate Change Initiative (ESA Cloud_cci) data: AVHRR-PM CLD_PRODUCTS
 v2.0 via Deutscher Wetterdienst.
- Stevens, B., M. Giorgetta, M. Esch, T. Mauritsen, T. Crueger, S. Rast, M. Salzmann, H. Schmidt,
 J. Bader, K. Block, R. Brokopf, I. Fast, S. Kinne, L. Kornblueh, U. Lohmann, R. Pincus, T.
 Reichler, and E. Roeckner (2013), Atmospheric component of the MPI-M Earth System
 Model: ECHAM6, Journal of Advances in Modeling Earth Systems, 5, 146-172, doi:
 10.1002/jame.20015.
- Stroeve, J. C., M. C. Serreze, M. M. Holland, J. E. Kay, J. Malanik, and A. P. Barrett (2012),
 The Arctic's rapidly shrinking sea ice cover: A research synthesis, Clim Change, 110, 10051027, doi: 10.1007/s10584-011-0101-1.
- Su, B., A. Wang, G. Wang, Y. Wang, T. Jiang (2016), Spatiotemporal variations of soil moisture
 in the Tarim River basin, China, Int. J. Appl. Earth Observ. Geoinform., 48, 122-130, doi:
 10.1016/j.jag.2015.06.012.
- Sus, O., M. Jerg, C. Poulsen, G. Thomas, S. Stapelberg, G. Mcgarragh, A. Povey, C. Schlundt,
 M. Stengel, and R. Hollmann (2016), The Community Cloud retrieval for CLimate (CC4CL).
 Part I: A framework applied to multiple satellite imaging sensors, manuscript in reparation.
- Susskind, J., C. Barnet, J. Blaisdell, L. Iredell, F. Keita, L. Kouvaris, G. Molnar, and M. Chahine
 (2006), Accuracy of geophysical parameters derived from Atmospheric Infrared
 Sounder/Advanced Microwave Sounding Unit as a function of fractional cloud cover, J.
 Geophys. Res., 111, D9, doi: 10.1029/2005JD006272.
- Tans, P., and R. Keeling (2015), http://www.esrl.noaa.gov/gmd/ccgg/trends/ (last accessed 31
 August 2016).
- Taylor, K. E. (2001), Summarizing multiple aspects of model performance in a single diagram. J.
 Geophys. Res., 106 (D7), 7183-7192.
- Taylor, K. E., R. Stouffer, and G. A. Meehl (2012), An overview of CMIP5 and the experimentdesign, Bull. Amer. Meteor. Soc., 93, 485-498.
- Teixeira, J., D. Waliser, R. Ferraro, P. Gleckler, T. Lee, and G. Potter (2014), Satellite
 Observations for CMIP5: The Genesis of Obs4MIPs, Bull. Amer. Meteor. Soc, 95, 13291334.

- Tian, B., E. J. Fetzer, B. H. Kahn, J. Teixeira, E. Manning, and T. Hearty (2013), Evaluating
 CMIP5 models using AIRS tropospheric air temperature and specific humidity climatology, J.
 Geophys. Res., 118, D50117, 114-134.
- Tsendbazar, N.-E., S. de Bruin, S. Fritz, and M. Herold (2015), Spatial Accuracy Assessment
 and Integration of Global Land Cover Datasets, Remote Sens, 7, 15804-15821.
- Van Roozendael, M., et al. (2015), ESA Ozone Climate Change Initiative (ESA Ozone_cci) data:
 L3S-TC-MERGED-DLR v1.0 via Centre for Environmental Data Analysis, 2 August 2013.
- Voldoire, A., E. Sanchez-Gomez, D. Salas y Mélia, B. Decharme, C. Cassou, S.Sénési, S.
 Valcke, I. Beau, A. Alias, M. Chevallier, M. Déqué, J. Deshayes, H. Douville, E. Fernandez,
 G. Madec, E. Maisonnave, M.-P. Moine, S. Planton, D.Saint-Martin, S. Szopa, S. Tyteca, R.
 Alkama, S. Belamari, A. Braun, L. Coquart, and F. Chauvin (2013), The CNRM-CM5.1
 global climate model: description and basic evaluation, Clim. Dyn., 40, 2091, doi:
 10.1007/s00382-011-1259-y.
- Volodin, E. M., N. A. Dianskii, and A. V. Gusev (2010), Climate model INMCM4.0, Izvestia
 RAS, Atmospheric and Oceanic Physics, 46(4), 448-466, doi: 10.1134/S000143381004002X.
- Wagner, W., W. Dorigo, R. de Jeu, D. Fernandez, J. Benveniste, E. Haas, M. Ertl (2012), Fusion
 of active and passive microwave observations to create an Essential Climate Variable data
 record on soil moisture, ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial
 Information Sciences, Volume I-7, 2012, XXII ISPRS Congress, 25 August 01 September
 2012, 315-231, Melbourne, Australia.
- Walsh, J. E., W. L. Chapman, and F. Fetterer (2015), Gridded Monthly Sea Ice Extent and
 Concentration, 1850 Onward, Version 1, Boulder, Colorado, USA.
- Wang, S., X. Mo, S. Liu, Z. Lin, and S. Hu (2016), Validation and trend analysis of ECV soil
 moisture data on cropland in North China Plain during 1981-2010, Int. J. Appl. Earth Observ.
 Geoinform., 48, 110-121, doi: 10.1016/j.jag.2015.10.010.
- Watanabe, M., T. Suzuki, R. O'ishi, Y. Komuro, S. Watanabe, S. Emori, T. Takemura, M.
 Chikira, T. Ogura, M. Sekiguchi, K. Takata, D. Yamazaki, T. Yokohata, T. Nozawa, H.
 Hasumi, H. Tatebe, and M. Kimoto (2010), Improved climate simulation by MIROC5: Mean
 states, variability, and climate sensitivity, J. Climate, 23, 6312-6335.
- Watanabe, S., T. Hajima, K. Sudo, T. Nagashima, T. Takemura, H. Okajima, T. Nozawa, H.
 Kawase, M. Abe, T. Yokohata, T. Ise, H. Sato, E. Kato, K. Takata, S. Emori, and M.
 Kawamiya (2011), MIROC-ESM 2010: model description and basic results of CMIP5-20c3m
 experiments, Geosci. Model Dev., 4, 845-872.
- Willet, K. M., A. J. Dolman, D. F. Hurst, J. Rennie, and P. W. Thorne (2014), [Global Climate]
 Overview [in: "State of the Climate 2013'], Bull. Am. Meteor. Soc., 95(7), S5-S8.

- Wise, M., K. Calvin, A. Thomson, L. Clarke, B. Bond-Lamberty, R. Sands, S. J. Smith, A.
 Janetos, and J. Edmonds (2009), Implications of limiting CO₂ concentrations for land use and
 energy, Science, 324, 1183-1186.
- 1698 World Meteorological Organization (WMO) (2014), Scientific Assessment of Ozone Depletion:
- 2014, World Meteorological Organization, Global Ozone Research and Monitoring Project Report No. 55, 416 pp., Geneva, Switzerland.
- Wu, T.-W. (2012), A mass-flux cumulus parameterization scheme for large-scale models:
 Description and test with observations, Clim. Dyn., 38, 725-744.
- Wu, T., R. Yu, F. Zhang, Z. Wang, M. Dong, L. Wang, X. Jin, D. Chen, L. Li (2010), The
 Beijing Climate Center for Atmospheric General Circulation Model (BCC-AGCM2.0.1):
 Description and its performance for the present-day climate, Clim. Dyn., 34, 123-147.
- Wunch, D., G. Toon, J. Blavier, R. Waschenfelder, J. Notholt, B. Connor, D. Griffith, and V.
 SherlockSherlock (2011), The Total Carbon Column Observing Network, Philos. T. Roy. Soc.
 A, 369, 2087-2112, 2011, doi: 10.1098/rsta.2010.0240.
- Wunch, D., G. C. Toon, V. Sherlock, N. M. Deutscher, X. Liu, D. G. Feist, and P. O. Wennberg
 (2015), The Total Carbon Column Observing Network's GGG2014 Data Version. Carbon
 Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge,
 Tennessee, USA, available at: doi: 10.14291/tccon.ggg2014.documentation.R0/1221662 (last
 accessed 27 November 2015).
- 1714 Yukimoto, S., H. Yoshimura, M. Hosaka, T. Sakami, H. Tsujino, M. Hirabara, T. Y. Tanaka, M.
- 1715 Deushi, A. Obata, H. Nakano, Y. Adachi, E. Shindo, S. Yabu, T. Ose, and A. Kitoh (2011),
- 1716 Meteorological Research Institute-Earth System Model Version 1 (MRI-ESM1) Model
- 1717 Description, Technical Report of the Meteorological Research Institute, 64, 83pp.

1718 Appendix – list of abbreviations and acronyms

1719	AATSR	Advanced Along-Track Scanning Radiometer	
1720	ACE	Atmospheric Chemistry Experiment	
1721	ADV	Advanced along-track scanning radiometer (AATSR) Dual-View	
1722	AEROCOM	Aerosol Comparisons between Observations and Models	
1723	AERONET	AErosol RObotic NETwork	
1724	AIRS	Atmospheric Infrared Sounder	
1725	AMSR-E	Advanced Microwave Scanning Radiometer - Earth Observing System	
1726	ana4MIPs	analyses for Model Intercomparison Projects	
1727	AOD	Aerosol Optical Depth	
1728	ATBD	Algorithm Theoretical Basis Documents	
1729	ATSR(-2)	Along-Track Scanning Radiometers (2)	
1730	AVHRR	Advanced Very High Resolution Radiometer	
1731	BDBP	Binary Data Base of Profiles	
1732	CALIOP	Cloud-Aerosol Lidar with Orthogonal Polarization	
1733	CC4CL	Community Cloud retrieval for CLimate	
1734	CCI	Climate Change Initiative	
1735	CERES	Clouds and the Earth's Radiant Energy System	
1736	CLARA-A2	CLoud, Albedo and RAdiation dataset, AVHRR-based	
1737	CMIP5/6	Coupled Model Intercomparison Project Phase 5/6	
1738	CMOR	Climate Model Output Rewriter	
1739	CMUG	Climate Modelling User Group	
1740	CO_2	carbon dioxide	
1741	CRESCENDO	Coordinated Research in Earth Systems and Climate: Experiments,	
1742	kNowledge, Dissemination and Outreach		
1743	CVDP	Climate Variability Diagnostics Package	
1744	DECK	Diagnostic, Evaluation and Characterization of Klima	
1745	DGVM	Dynamic Global Vegetation Model	
1746	DJF	December, January, February	
1747	DU	Dobson Unit	
1748	EBAF	Energy Balanced And Filled	
1749	ECV	Essential Climate Variable	
1750	ENVISAT	Environmental Satellite	
1751	ENSO	El Niño Southern Oscillation	
1752	ERS-2	European Remote Sensing Satellite 2	
1753	ESA	European Space Agency	
1754	ESGF	Earth System Grid Federation	
1755	ESM	Earth System Model	
1756	ESMValTool	Earth System Model Evaluation Tool	
1757	EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites	

1758	FMI	Finnish Meteorological Institute
1759	FTS	Fourier Transform Spectrometer
1760	GCOS	Global Climate Observing System
1761	GHG	greenhouse gases
1762	GOME/-2	Global Ozone Monitoring Experiment / 2
1763	GOMOS	Global Ozone Monitoring by Occultation of Stars
1764	GOSAT	Greenhouse gases observing satellite
1765	GPCP	Global Precipitation Climatology Project
1766	GRACE	Gravity Recovery and Climate Experiment
1767	HadISST	Hadley Centre Sea Ice and Sea Surface Temperature data set
1768	IASI	Infrared Atmospheric Sounding Interferometer
1769	IGAC	International Global Atmospheric Chemistry
1770	ITCZ	Inter-Tropical Convergence Zone
1771	JAG	International Journal of Applied Earth Observation and Geoinformation
1772	JJA	June, July, August
1773	L2/3/4	Level 2/3/4
1774	MAN	Marine Aerosol Network
1775	MERIS	MEdium Resolution Imaging Spectrometer
1776	Metop	Meteorological Operational Satellite
1777	MIPAS	Michelson Interferometer for Passive Atmospheric Sounding
1778	MMM	Multi-Model Mean
1779	MODIS	Moderate Resolution Imaging Spectroradiometer
1780	NASA	National Aeronautics and Space Administration
1781	NCAR	National Center for Atmospheric Research
1782	NCEP	National Centers for Environmental Prediction
1783	NDVI	Normalized Differenced Vegetation Index
1784	NH	Northern Hemisphere
1785	NIR	near-infrared
1786	NIWA	National Institute of Water and Atmospheric Research
1787	NOAA	National Oceanic and Atmospheric Administration
1788	NSIDC	National Snow and Ice Data Center
1789	obs4MIPs	observations for Model Intercomparison Projects
1790	OMI	Ozone Monitoring Instrument
1791	ORAC	Oxford-Rutherford Appleton Laboratory (RAL) Aerosol and Clouds
1792	OSI SAF	Satellite Application Facility on Ocean and Sea Ice
1793	OSIRIS	Optical Spectrograph and InfraRed Imaging System
1794	OSTIA	Operational Sea surface Temperature and sea-Ice Analysis
1795	PATMOS-x	Pathfinder Atmospheres Extended
1796	PCMDI	Program for Climate Model Diagnostics and Intercomparison
1797	PFT	Plant Functional Type

1798	POLDER	POLarization and Directionality of the Earth's Reflectances
1799	PMP	PCMDI metrics package
1800	RAL	Rutherford Appleton Laboratory
1801	RCP	Representative Concentration Pathways
1802	RMSD	relative space-time Root-Mean-Square Deviation
1803	RMSE	Root-Mean-Square Error
1804	SCIAMACHY	Scanning Imaging Absorption Spectrometer for Atmospheric CHartographY
1805	SECM	Simple Empirical CO ₂ Model
1806	SH	Southern Hemisphere
1807	SI	Sea Ice
1808	SMR	Sub-Millimetre Radiometer
1809	SPARC	Stratospheric Processes and their Role in Climate
1810	SPOT	Satellite Pour l'Observation de la Terre
1811	SSM/I	Special Sensor Microwave Imager
1812	SWIR	short-wave infrared
1813	SST	Sea Surface Temperature
1814	SU	Swansea University retrieval algorithm
1815	TANSO	Thermal And Near-infrared Sensor for carbon Observation
1816	TCCON	Total Carbon Column Observation Network
1817	UV	ultraviolet
1818	Vis	visible spectral range
1819	WCRP	World Climate Research Programme
1820	WMO	World Meteorological Organization